Check for updates

Received: 16 August 2023 | Accepted: 21 August 2023

DOI: 10.1111/jsr.14035

REVIEW ARTICLE

The European Insomnia Guideline: An update on the diagnosis and treatment of insomnia 2023

Dieter Riemann[1,2] | Colin A. Espie[3] | Ellemarije Altena[4] | Erna Sif Arnardottir[5,6] | Chiara Baglioni[7] | Claudio L. A. Bassetti[8] | Celyne Bastien[9] | Natalija Berzina[10] | Bjørn Bjorvatn[11] | Dimitris Dikeos[12] | Leja Dolenc Groselj[13] | Jason G. Ellis[14] | Diego Garcia-Borreguero[15] | Pierre A. Geoffroy[16] | Michaela Gjerstad[17] | Marta Gonçalves[18] | Elisabeth Hertenstein[19] | Kerstin Hoedlmoser[20] | Tuuliki Hion[21] | Brigitte Holzinger[22] | Karolina Janku[23] | Markus Jansson-Fröjmark[24,25] | Heli Järnefelt[26] | Susanna Jernelöv[24,25] | Poul Jørgen Jennum[27] | Samson Khachatryan[28] | Lukas Krone[3,8,19] | Simon D. Kyle[3] | Jaap Lancee[29] | Damien Leger[30] | Adrian Lupusor[31] | Daniel Ruivo Marques[32,33] | Christoph Nissen[34] | Laura Palagini[35] | Tiina Paunio[36] | Lampros Perogamvros[34] | Dirk Pevernagie[37] | Manuel Schabus[20] | Tamar Shochat[38] | Andras Szentkiralyi[39] | Eus Van Someren[40,41] | Annemieke van Straten[42] | Adam Wichniak[43] | Johan Verbraecken[44] | Kai Spiegelhalder[1]

Correspondence
Dieter Riemann, Department of Clinical Psychology and Psychophysiology, Centre for Mental Disorders, Medical Centre – University of Freiburg, Faculty of Medicine, University of Freiburg, Germany; Hauptstr. 5, D-79104 Freiburg, Germany.
Email: dieter.riemann@uniklinik-freiburg.de

Summary

Progress in the field of insomnia since 2017 necessitated this update of the European Insomnia Guideline. Recommendations for the diagnostic procedure for insomnia and its comorbidities are: clinical interview (encompassing sleep and medical history); the use of sleep questionnaires and diaries (and physical examination and additional measures where indicated) (A). Actigraphy is not recommended for the routine evaluation of insomnia (C), but may be useful for differential-diagnostic purposes (A). Polysomnography should be used to evaluate other sleep disorders if suspected (i.e. periodic limb movement disorder, sleep-related breathing disorders, etc.), treatment-resistant insomnia (A) and for other indications (B). Cognitive-behavioural therapy for insomnia is recommended as the first-line treatment for chronic insomnia in adults of any age (including patients with comorbidities), either applied in-person or digitally (A). When cognitive-behavioural therapy for insomnia is not sufficiently

For affiliations refer to page 26

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.
© 2023 The Authors. *Journal of Sleep Research* published by John Wiley & Sons Ltd on behalf of European Sleep Research Society.

effective, a pharmacological intervention can be offered (A). Benzodiazepines (A), benzodiazepine receptor agonists (A), daridorexant (A) and low-dose sedating antidepressants (B) can be used for the short-term treatment of insomnia (≤ 4 weeks). Longer-term treatment with these substances may be initiated in some cases, considering advantages and disadvantages (B). Orexin receptor antagonists can be used for periods of up to 3 months or longer in some cases (A). Prolonged-release melatonin can be used for up to 3 months in patients ≥ 55 years (B). Antihistaminergic drugs, antipsychotics, fast-release melatonin, ramelteon and phytotherapeutics are not recommended for insomnia treatment (A). Light therapy and exercise interventions may be useful as adjunct therapies to cognitive-behavioural therapy for insomnia (B).

KEYWORDS
diagnosis, evidence-based medicine, guideline, insomnia, treatment

1 | SUMMARY FOR PATIENTS

1.1 | What is insomnia?

Insomnia is a sleep disorder where people struggle to get off to sleep or to stay asleep. Some individuals have both issues, and others may also have early-morning awakenings, where they are unable to get back to sleep after awakening earlier than desired. Importantly, these night-time sleep difficulties are coupled with significant daytime problems that affect the person's ability to function at their best. Daytime fatigue, low mood or irritability, and problems with attention or concentration are usually experienced. To be diagnosed with an "insomnia disorder", these difficulties have to occur at least several times a week over a period of 3 months. Insomnia is a very common disorder (up to 10% of the adult population in Europe) and, in addition to a great deal of personal suffering, it also results in increased costs to healthcare services and to society at large (e.g. reduced productivity at work).

1.2 | How can insomnia be treated?

Currently, there are two ways to treat insomnia. According to scientific evidence the first, and the most effective, approach is cognitive-behavioural therapy for insomnia (CBT-I). As the name suggests, CBT-I addresses the mental or cognitive aspects of insomnia (e.g. the racing mind), and the behavioural aspect reestablishes a healthy sleep pattern. CBT-I can be offered as single or group therapy (face-to-face [F2F]) or as a digital therapy, where it is delivered as a web-based intervention or on a treatment-based app. The second approach to treat insomnia is pharmacological (i.e. pill-based). There is a variety of sleep medications available, but it is recommended that these are only taken for a short period (no longer than 4 weeks) to avoid the body getting used to them or becoming dependent upon them. In some cases, after weighing the advantages and disadvantages, some medications may be given for longer periods of time.

1.3 | Who developed this guideline?

This European Insomnia Guideline 2023 was developed by a group of researchers and clinicians in the European Sleep Research Society (ESRS), and the European Insomnia Network (EIN).

1.4 | Which treatment is recommended by this guideline?

It is recommended that all patients with insomnia, whether they have other medical conditions or mental health problems, or not, are offered CBT-I as their initial treatment. CBT-I may be delivered by a clinician or therapist (F2F), or (preferably guided) digitally using a scientifically demonstrated web or mobile treatment platform. If this approach is not sufficiently effective, it is recommended that patients and their treating physicians should come to a shared decision about whether or not medication might be initiated. At present, the evidence suggests that drug treatments in general should be limited to, at most, 4 weeks in duration, and even then with care: tolerance develops within days to weeks. Dose increases are not advised, and may accelerate the development of dependence. In some cases, longer treatment periods may be indicated, carefully weighing the advantages and disadvantages.

2 | GUIDELINE REPORT

This guideline is an update of the European Insomnia Guideline that was published in 2017 (Riemann, Baum, et al., 2017), and developed by a task force of the ESRS and the EIN. The European Insomnia Guideline was based on the German Insomnia Guideline (Riemann, Baglioni, et al., 2017) and was endorsed by the World Sleep Society (Morin et al., 2021). A revision of the German Insomnia Guideline is underway (Spiegelhalder et al., 2023). This first update of the European Insomnia Guideline is inspired by and draws upon this revision of the German Insomnia Guideline.

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

This guideline focuses on the target population of adult patients suffering from chronic insomnia as defined by the International Classification of Diseases (ICD-10/ICD-11). This includes all subtypes of insomnia, for example, non-organic insomnia/chronic insomnia and insomnia comorbid with somatic (formerly named "organic" insomnia) or mental disorders. The guideline addresses adult patients over the age of 18 years. The literature on insomnia in children and adolescents was not reviewed.

The guideline is based on a review of all relevant international literature and has a particular salience to the provision of clinical services across Europe. It will be of interest to health professionals who are involved in the diagnosis and treatment of insomnia on either an out- or in-patient basis. Most insomnia is managed by general practitioners in primary care settings, and by clinicians who are not specialists in sleep medicine. The guideline should also be helpful to such individuals (i.e. who treat insomnia in routine clinical care without access to advanced expertise or facilities). The guideline should also be useful to specialists in psychiatry, clinical psychology/psychotherapy, psychosomatic medicine, neurology, occupational medicine, pharmacy/pharmacology and other medical specialties who commonly see patients with insomnia in the context of other comorbid physical and mental health conditions. Finally, the guideline will be especially relevant to professionals trained in sleep medicine and who are members of, or are credentialled by, the ESRS.

The revised guideline highlights aspects of clinical management that reflect advances in knowledge and practice that can be delineated from the updated evidence. Accordingly, less emphasis is placed upon detailed reproduction of extant information that is already outlined in the 2017 version (Riemann, Baglioni, et al., 2017).

2.1 | Literature search

The 2023 update was designed to build upon scientific knowledge and clinical recommendations from the first guideline, which covered evidence up to June 2016 (Riemann, Baglioni, et al., 2017; Riemann, Baum, et al., 2017). This strategy therefore aimed to both complement and extend previous literature searches, while applying a consistent methodology.

Therefore, to identify relevant studies on the topic of insomnia, a systematic literature search (English language articles only) was conducted using the databases Pubmed and Cochrane Library (www.cochranelibrary.com) for the period from June 2016 until October 2022 (with a further update till May 2023 added). For this update, primarily meta-analyses were identified as the basis for grading recommendations. If there were several meta-analyses on a given topic, the most recent and qualitatively better meta-analyses were chosen to be presented in the first instance. The quality of meta-analyses was judged by methodological rigour, like low risk of bias, number of included studies or sample sizes. For topics without published meta-analyses, systematic reviews or qualitatively adequate randomised controlled studies were used.

The following keywords were used for literature search:

For the area of **non-pharmacological treatments**, the keyword "insomnia" was searched in connection with other keywords: "sleep hygiene", "relaxation", "mindfulness", "behavior therapy", "cognitive therapy", "cognitive behavioral therapy", "stimulus control", "sleep restriction", "psychotherapy", "light therapy", "exercise", "music", "non-invasive brain stimulation".

For the area of **pharmacological treatments**, the keyword "insomnia" was searched in connection with other keywords: "benzodiazepine", "benzodiazepine receptor agonist", "sedating antidepressant", "antipsychotic", "neuroleptic", "orexin", "antihistaminic", "herbal", "phytotherapy", "melatonin".

Furthermore, the journal *Sleep Medicine Reviews* was hand searched for meta-analyses on the diagnosis and treatment of insomnia. All issues of this journal until October 2022 were incorporated, furthermore articles were incorporated that were "in press".

2.2 | Writing and consensus

The first draft of this update was formulated and written by Dieter Riemann and Kai Spiegelhalder, following partly, and where adequate, the update of the German insomnia guideline (Spiegelhalder et al., 2023). In the next step, all involved authors received the first draft of the guideline (15 April 2023) and were asked to provide feedback within a period of 4 weeks (15 May 2023). After receiving feedback and incorporating suggested changes/improvements in the second draft of the guideline, two online meetings were held on 21 July and 1 August 2023 to discuss this version of the guideline and to reach consensus. A third draft of the guideline was then sent out asking for final consent from all authors by 7 August 2023. Finally, the guideline was approved by the guideline committee of the ESRS and by the ESRS board, before submission to *Journal of Sleep Research* (JSR).

2.3 | Grading of the evidence/recommendations

In order to grade the evidence of included studies/meta-analyses to update the recommendations in the guideline, a procedure similar to that already outlined in 2017 (Lorenz et al., 2001; Riemann, Baglioni, et al., 2017; Riemann, Baum, et al., 2017) was followed (Table 1).

The transformation of grades of evidence into grades of recommendations was performed according to this scheme and through consensus decision between all involved authors. For more details, see supplemental material in Riemann, Baum, et al. (2017). Instead of only using two types of recommendations (strong versus weak; Riemann, Baum, et al., 2017), we used four steps of recommendations for this update, ranging from A (very strong recommendation), B (strong),

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

C (weak) to D (very weak recommendation). Please note; levels of evidence do not always directly translate into grades of recommendation, as suggested in Table 1; in some cases, a consensus decision became the decisive factor for the grading of the recommendation.

Reported effect sizes from the meta-analyses were graded as follows: effect sizes (Cohen's *D*) ≥ 0.2– < 0.5: small effect; effect sizes ≥ 0.5– < 0.8: medium effect; effect sizes ≥ 0.8: large effect.

3 | DIAGNOSIS OF INSOMNIA

The 2017 version of this guideline reflected upon marked advances in understanding insomnia as a disorder in its own right (insomnia disorder), that had been recommended by the DSM-5 (Diagnostic and Statistical Manual of The American Psychiatric Association; APA, 2013) and the third edition of ICSD-3 (International Classification of Sleep Disorders; AASM, 2014). Practicing clinicians in most European countries, however, generally have to adhere to the International Classification of Diseases (ICD: World Health Organisation). It is noteworthy, therefore, that the latest revision (ICD-11) was endorsed by the World Health Assembly at its 72nd meeting in 2019 and came into effect globally on 1 January 2022 (WHO, 2022). Whereas the previous version (ICD-10; WHO, 1994) differentiated "non-organic" versus "organic insomnia", this distinction was abandoned as not evidence based in ICD-11 in favour of the comprehensive category "chronic insomnia" (Code 7A00; and "short-term insomnia" for "transient insomnia"). Criteria for insomnia disorder according to DSM-5, ICSD-3 and ICD-11 are now broadly aligned, and are depicted in Tables 2 and 3.

TABLE 1 Study types, hierarchy of evidence and grades of recommendation

Type of study	Level of evidence	Grade of recommendation
Systematic review/meta-analysis of RCTs	1a	A
One high-quality RCT	1b	A
"All or nothing principle"	1c	B
Systematic review of high-quality cohort studies	2a	B
One cohort study/RCT with adequate design but moderate data quality	2b	B
Outcome research studies	2c	B
One systematic review of case control studies	3a	B
One case control study	3b	B
Case series/cohort or case control studies of moderate quality	4	C
Expert opinion, etc.	5	D

Abbreviation: RCT, randomised–controlled study.

TABLE 2 Diagnostic criteria for chronic insomnia disorder according to ICSD-3 (AASM, 2014)

A. The patient reports, or the patient's parent or caregiver observes, one or more of the following:
   1. Difficulty initiating sleep
   2. Difficulty maintaining sleep
   3. Waking up earlier than desired
   4. Resistance to going to bed on appropriate schedule
   5. Difficulty sleeping without parent or caregiver intervention

B. The patient reports, or the patient's parent or caregiver observes, one or more of the following related to the nighttime sleep difficulty:
   1. Fatigue/malaise
   2. Attention, concentration or memory impairment
   3. Impaired social, family, occupational or academic performance
   4. Mood disturbance/irritability
   5. Daytime sleepiness
   6. Behavioural problems (e.g. hyperactivity, impulsivity, aggression)
   7. Reduced motivation/energy/initiative
   8. Proneness for errors/accidents
   9. Concerns about or dissatisfaction with sleep

C. The reported sleep/wake complaints cannot be explained purely by inadequate opportunity (i.e. enough time is allotted for sleep) or inadequate circumstances (i.e. the environment is safe, dark, quiet and comfortable) for sleep

D. The sleep disturbance and associated daytime symptoms occur at least three times per week

E. The sleep disturbance and associated daytime symptoms have been present for at least 3 months

F. The sleep/wake difficulty is not better explained by another sleep disorder

TABLE 3 Diagnostic criteria for chronic insomnia according to ICD-11 (Code 7A00)

(https://icd.who.int/browse11/l-m/en#/http://id.who.int/icd/entity/323148092)

Chronic insomnia is a frequent and persistent difficulty initiating or maintaining sleep that occurs despite adequate opportunity and circumstances for sleep, and that results in general sleep dissatisfaction and some form of daytime impairment. Daytime symptoms typically include fatigue, depressed mood or irritability, general malaise, and cognitive impairment. The sleep disturbance and associated daytime symptoms occur at least several times per week for at least 3 months. Some individuals with chronic insomnia may show a more episodic course, with recurrent episodes of sleep/wake difficulties lasting several weeks at a time over several years. Individuals who report sleep-related symptoms in the absence of daytime impairment are not regarded as having an insomnia disorder. If the insomnia is due to another sleep–wake disorder, a mental disorder, another medical condition, or a substance or medication, chronic insomnia should only be diagnosed if the insomnia is an independent focus of clinical attention.

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

In this update we will not comment on aetiological and/or pathophysiological concepts of insomnia, as this would go far beyond the remit of this guideline. The interested reader is referred to recent work, much of which is part of the special insomnia issue of JSR (Dressle & Riemann, 2023; Espie, 2023; Fernandez & Perlis, 2023; Palagini et al., 2023; Reffi et al., 2023; Riemann et al., 2022; Tang et al., 2023; van Someren, 2021).

An overview of the relevant diagnostic procedures is given in Table 4.

3.1 | Insomnia complaint

The diagnosis of chronic insomnia disorder according to DSM 5, ICSD-3 and ICD-11 continues to be solely based on taking a clinical history (anamnesis) and the patient's self-reports of sleep-onset or sleep maintenance difficulties, or early-morning awakening. Likewise, clinical consideration of whether or not these sleep symptoms are coupled with dissatisfaction with sleep and any attributed daytime impairment is mandatory, as well as that symptoms occur despite adequate time allotted for sleep and the opportunity to sleep in a comfortable environment. Quantitative criteria related to sleep-onset latency, sleep duration or the frequency of nocturnal awakenings do not have to be fulfilled in order to diagnose insomnia disorder. The complaint of insomnia occurring at least three nights per week for 3 months with associated daytime sequelae represents minimal criteria for this clinical evaluation of chronic insomnia disorder.

TABLE 4 Diagnostic management of insomnia with or without comorbidities

General anamnesis and examination (A)
• Former and present somatic, neurological and mental disorders
• Personality factors, work and partnership situation, interpersonal conflicts
• Substance use (medication, alcohol, caffeine, nicotine, illegal drugs)
• Physical examination (if indicated)
• Additional measures (if indicated): laboratory testing including, e.g. blood count, thyroid, hepatic and renal parameters, C-reactive protein, haemoglobin, ferritin and vitamin B12; if indicated: electrocardiogram, electroencephalogram, computed tomography/Magnetic resononance tomography, circadian markers (melatonin, core body temperature)
• The anamnesis should include caregivers if necessary
Sleep history (A)
• History of the sleep complaints and daytime functioning
• Information from bed partner/caregivers (snoring, breathing pauses = apneas, periodic limb movements during sleep, nocturnal restlessness, "strange" behaviours)
• Work time/circadian factors (shift- and night work, phase advance, delay)
• Sleep–wake pattern, including daytime sleep (sleep diary, sleep questionnaires)
Actigraphy
• In case of clinical suspicion of irregular sleep–wake schedules or circadian rhythm sleep–wake disorders (A)
• In case of clinical suspicion of periodic leg movements in sleep (A)
• To assess quantitative rest activity (A) and sleep parameters (C)
PSG
• In case of clinical suspicion of comorbid sleep disorders (A)
• Treatment-resistant insomnia (A)
• In case of clinical suspicion of large discrepancy between subjectively experienced and PSG-measured sleep (B)

Abbreviation: PSG, polysomnography.

3.2 | Sleep diaries and questionnaires

As recognised in the 2017 guideline (Riemann, Baglioni, et al., 2017), the evaluation of insomnia should be supported by the use of a sleep diary for a period of at least 7–14 days (see consensus sleep diary: Carney et al., 2012). The Insomnia Severity Index (ISI; Bastien et al., 2001) is an established tool to gauge the severity of the insomnia. The ISI can range with a score from 0 to 28. Values from 8 to 14 suggest subclinical insomnia, values from 15 to 21 moderately severe insomnia, and values from 22 to 28 severe insomnia. The authors recommend a cut-off score of 10 for caseness in community settings, and a change score of 8.4 as a sign of moderate improvement (Morin et al., 2011). The ISI has a two-item version that can be used for weekly monitoring or other situations where a shorter version may be needed (Kraepelien et al., 2021) The eight-item Sleep Condition Indicator (SCI), based upon DSM-5 criteria for insomnia disorder, is an alternative instrument to the ISI (Espie, Kyle, Hames, et al., 2014). The SCI also has a two-item short-form version (Luik et al., 2019), and has age and sex reference data on a sample of 200,000 adults (Espie et al., 2018), with a Reliable Change Index of 7 scale points. The frequently used Pittsburgh Sleep Quality Index (PSQI; Buysse et al., 1989) may be helpful to gain a broader picture of sleep-related symptoms that can be relevant for the judgement of insomnia. The PSQI delivers values from 0 to 21, with values above 5 supposedly reflecting clinically relevant sleep impairments. It should be noted, however, that the PSQI is not a measure of insomnia disorder, but it broadly asks about symptoms from a range of other sleep disorders.

In the meantime, a variety of questionnaires has become available, not only to measure insomnia, but for many other aspects of other sleep disorders (the interested reader is referred to Shahid et al., 2012). A single comprehensive and validated questionnaire for sleep and daytime impairment for clinical use has not been developed yet, to our knowledge, and is urgently needed.

3.3 | Actigraphy

Although the literature on actigraphy continues to expand, and its availability may become more pervasive, there is limited support for the mandatory application of actigraphy in the routine clinical

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

evaluation and diagnosis of insomnia. Medical grade actigraphic devices may be used to evaluate "free living" bedtimes and sleep times, and their variability, at a rather low cost for longer periods of time, and so have particular application where a circadian rhythm sleep–wake disorder is suspected (Smith et al., 2018). According to two systematic reviews, many devices based on actigraphy underestimate the severity of sleep disorder symptoms and overestimate sleep duration compared with polysomnography (PSG; Kolla et al., 2016; Smith et al., 2018). Rösler et al. (2023) concluded that actigraphy was not able to differentiate patients with insomnia from good sleepers. Actigraphy may be primarily helpful in a differential-diagnosis, to identify irregular bedtime patterns—actigraphy, however, does not deliver a valid estimation of sleep stages like the PSG.

Further, there are numerous lifestyle products available, in the form of watches or similar wearable devices, that incorporate some form of actigraphic and/or heart rate measurement. However, the validity and utility of such devices for the diagnosis of insomnia has not been satisfactorily researched or demonstrated at this point in time (Kolla et al., 2016). It has even been suggested that using these devices might have a negative impact on sleep behaviour.

3.4 | Polysomnography (PSG)

It is well established that PSG is indicated when there is clinical suspicion of other sleep disorders like periodic limb movement disorder (PLMD) or any kind of sleep-related breathing disorders (SRBD; Crönlein et al., 2012), such as obstructive sleep apnea (OSA). Because insomnia does not protect against, or exclude, such disorders, PSG is equally indicated if clinical suspicion is present in people with insomnia. However, PSG is not necessary or sufficient for the diagnosis of insomnia per se (Dikeos et al., 2023). A meta-analysis of the PSG literature in insomnia showed that, as compared with controls without sleep complaints, patients with insomnia have a significant reduction in sleep efficiency, a reduction in total sleep time, an increase in nocturnal wake times, and reductions in slow-wave sleep and rapid eye movement (REM) sleep (Baglioni et al., 2014). These differences compared with good sleepers, however, are considerably lower than would be expected from subjective (i.e. sleep diaries/questionnaires) measures of sleep. Moreover, similar deviations can be found in several other sleep disorders and other conditions, resulting in insufficient specificity. Although it has been suggested that feedback about any discrepancy between physiological and objective and subjective measures might be used therapeutically (Tang & Harvey, 2006), CBT-I and all other treatments are substantially effective for insomnia disorder in the absence of such data. We recommend that PSG should also be performed in patients with insomnia who have been refractory to various therapeutic interventions (CBT-I and/or hypnotics) with the aim to detect hitherto occult sleep disorders, in patients at risk for tiredness/fatigue-related accidents (i.e. professional drivers, etc.: to gauge the extent of sleep deprivation) and in patients where a huge mismatch between subjective and objective findings is suspected. Riemann et al. (2022) have outlined that a potential explanation for the discrepancy between subjective and objective findings may be via an altered sleep microstructure in insomnia, including an increased frequency of micro-arousals (especially in REM sleep) and increased amounts of fast-frequency waves (in the sigma and beta-band).

3.5 | Medical evaluation

Table 5 lists a wide range of medical, mental, neurological disorders and substances, which are implicated in this context (please note: probably any kind of medical disease might be involved as a causative or contributive factor for insomnia).

A thorough physical examination is therefore indicated in patients with insomnia with suspected comorbidities, to be able to exclude specific medical disorders that are in need of specific treatment and for, when necessary, treating insomnia as a comorbid condition. The historical and now outdated clinical perspective, that insomnia is primarily a symptom, was strongly challenged by evidence that chronic insomnia is an independent risk factor for mental health conditions such as depression or anxiety disorders (Baglioni et al., 2011; Hertenstein et al., 2019). Indeed, almost any kind of mental disorder is frequently accompanied by insomnia or can contribute to a worsening of insomnia symptoms (Table 5). Many patients with insomnia suffer comorbidly from another mental disorder, which may be stigmatised and therefore will not be reported spontaneously by patients. Therefore, within the context of the clinical evaluation of insomnia, any kind of mental disorder should be actively explored.

Insomnia is highly prevalent in such neurological disorders as stroke, Parkinson's disease, epilepsy, headache/migraine and traumatic brain injury (Bassetti et al., 2015, 2020). More recently, insomnia has also been shown to represent an independent risk factor for some neurological disorders such as stroke and dementia (Damsgard et al., 2022; Zheng et al., 2019).

Insomnia is highly prevalant also in medical disorders including diabetes (LeBlanc et al., 2018) and cardiovascular disease/hypertension (Li, Gan, et al., 2021; Li, Li, et al., 2021; Sofi et al., 2014). Indeed, it seems there is a truly bi-directional relationship between sleep disturbances and health conditions, such that insomnia should be evaluated and actively treated in comorbid conditions. Moreover, this underscores medical and psychiatric evaluations being part of a thorough insomnia assessment. Accordingly, normal sleep is recognised as an important predictor of brain and mental health (Bassetti et al., 2015).

Special note must be made of other sleep disorders, like SRBD/OSA, restless legs syndrome (RLS) and PLMD. Only recently has it become evident that there is a considerable percentage of patients with OSA also presenting with insomnia—now termed COMISA (comorbid insomnia and sleep apnea). Recent meta-analyses indicate that 30%–40% of patients with insomnia may have OSA, and 30%–50% of patients with OSA meet insomnia criteria (Zhang, Ren, Zhou, et al., 2019; Sweetman et al., 2021). These data argue for a strict inclusion of an OSA screening tool (e.g. STOP-Bang questionnaire; Chung et al., 2008) in the diagnostic process for insomnia and

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 5 Major medical comorbidities or contributory factors to chronic insomnia

Mental	Medical	Neurological	Substance use/dependence
Depressive disorders	Cardiovascular disorders	Neurodegenerative diseases	Alcohol
Bipolar disorders	Diabetes mellitus	Cerebrovascular diseases	Nicotine
Anxiety disorders	Chronic kidney diseases	Traumatic brain injury	Caffeine
Borderline personality disorder	Chronic obstructive pulmonary diseases	Multiple sclerosis	Tetrahydrocannabinol /marihuana
Posttraumatic stress disorder	Rheumatic disorders	RLS/PLMD	Opioids
Schizophrenia	Chronic pain	Fatal familial insomnia	"Designer" drugs
Substance use disorders	Any kind of malignant disorder		Cocaine
	SRBD/OSA		Amphetamines

Abbreviations: OSA, obstructive sleep apnea; PLMD, periodic limb movement disorder; RLS, restless legs syndrome; SRBD, sleep-related breathing disorder.

conversely insomnia screening for each patient with suspected SRBD. If the screening questionnaire is positive for OSA, a home sleep apnea test or PSG is required for a valid diagnosis of OSA. Insomnia is a crucial symptom of many patients with RLS. PLMD extremely frequently accompanies RLS and may contribute independently to the experience of insomnia. To determine the extent to which PLMD/RLS may be contributing to the insomnia, the International Restless Legs Syndrome Study Group Rating Scale (IRLSGR; Abetz et al., 2006) should be used to gauge the occurrence and severity of symptoms. The diagnosis of RLS is mainly a clinical diagnosis, but PLMD requires a PSG.

Given the high comorbidity between insomnia complaints and OSA/SRBD, PLMD and RLS, the diagnostic and differential-diagnostic process should pay utmost attention to delineate the probable bi-directional relationships between insomnia and the "other" sleep disorder—this information should also directly inform the therapeutic process, meaning that in many cases the insomnia and the "other" sleep disorder will be treated simultaneously according to suggested guidelines/articles (Garcia-Borreguero et al., 2016; Sweetman et al., 2023).

Furthermore, a number of substances may lead to the development, maintenance or aggravation of insomnia.

Alcohol consumption may play a prominent role in this context (Perney & Lehert, 2018; Hu et al., 2020; Hertenstein et al., 2019). Frequently, alcohol is also used mal-adaptively as self-medication to self-manage sleep-onset or sleep maintenance problems. Therefore, alcohol consumption and other substances should be specifically probed. Insomnia as a possible side-effect of prescribed medications may occur with a huge variety of substances, including corticosteroids, beta-blockers, beta-sympathomimetics, antibiotics, antidementives, selective serotonin reuptake blockers and probably many more (for an overview, see Riemann & Nissen, 2012; Schierenbeck et al., 2008) impacting on sleep.

It is important to know even in clear cases of the causative/maintaining role of a medical or mental disorder or consumption of a given substance, the typical psychophysiological vicious circle of insomnia may have developed in many patients. This includes anticipatory anxiety of a poor night, nocturnal ruminations, increased psychophysiological arousal and behavioural changes like excessive time in bed. Insofar, these clinical phenomena also may need to be addressed as part of the therapeutic approach towards comorbid insomnias.

Please note that with DSM-5, ICSD-3 and ICD-11, comorbidity of insomnia with any kind of medical disorder is acknowledged—the former hierarchy of primary versus secondary insomnia was abandoned because it lacked sufficient evidence, especially the assumed direct causality between medical disorders and associated insomnia symptoms, which should disappear when the primary disorder is treated, never became evidence-based.

In summary, the diagnostic and differential-diagnostic process for chronic insomnia disorder is complex and may require not only a clinician's direct time, but also the involvement of medical tests and interdisciplinary collaborations between different medical specialties. The process is not made easier due to the intricate bi-directional relationships between insomnia and almost any kind of medical disorder. Comorbidity may be more the rule than the exception, which entails the integration of insomnia treatments under a very broad medical context.

Needless to say, a proper diagnostic and differential diagnostic evaluation has to take place prior to any therapy, be it CBT-I, digital CBT-I (dCBT-I) or any kind of pharmacological treatment.

4 | TREATMENT OF INSOMNIA

Up to 2016/2017, published insomnia guidelines tended to favour pharmacotherapy over non-pharmacological approaches. Both fields existed in parallel worlds (i.e. on the one hand medical specialists favouring pharmacotherapy, and on the other hand, mainly clinical psychologists and psychotherapists, favouring CBT-I or other forms of psychotherapy). This antagonism seems to have been reconciled over the last decade (see guidelines/summaries by Quaseem et al., 2016; Wilson et al., 2019; Edinger et al., 2021), and is reflected by the wide authorship of this guideline.

One crucial question when discussing insomnia therapy is the question who and how many patients with insomnia seek help from

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

professional healthcare. Data from Germany (Marschall et al., 2017; data from health insurance) indicate that probably not more than 30% of those afflicted with insomnia have ever sought medical help. Considering over the last 12 months, only 15% indicated having seen a doctor specifically because of their insomnia. These sobering data need to be borne in mind when considering the treatment recommendations of this guideline. Given the bi-directional relationships between insomnia and all medical disorders, a strong case can be made to raise not only public awareness but also of the medical community concerning insomnia.

Concerning the choice of treatment option, we strongly suggest to follow a shared decision-making approach (Bomhof-Roordink et al., 2019). Diagnostic and treatment options should be outlined by the clinician and discussed together with the patient.

4.1 | Non-pharmacological treatments

4.1.1 | Cognitive-behavioural therapy for insomnia (CBT-I)

The CBT-I is typically provided as a multicomponent treatment comprising psychoeducation (including sleep hygiene), relaxation therapy, sleep-restriction therapy (SRT), stimulus control therapy (SCT) and several cognitive therapeutic strategies. These are administered over four–eight therapy sessions in single or group format by certified health professionals (mainly clinical psychologists/psychological or medical psychotherapists). Baglioni et al. (2022) summarised these treatments in a comprehensive manual/textbook aimed at health professionals involved in the treatment of insomnia. Chapter 2 in this book (pp. 19–41) describes the CBT-I "standard protocol", and provides detailed information on the scientific basis, therapy rationale and treatment instructions for each CBT-I component (Espie, 2022a). Baglioni, Altena, et al. (2020), have summarised CBT-I training strategies and the dissemination of CBT-I to health professionals. The website of the ESRS lists presently available teaching courses for health professionals in Europe (www.esrs.eu).

The multicomponent delivery approach is typical of the pattern of CBT provision in other common disorders, such as anxiety and depression. That said, substantial early evidence was gathered on single component treatments for insomnia (e.g. progressive relaxation, stimulus control, paradoxical intention, sleep restriction), and there has been a return to evaluating separate components in recent years. This is in keeping with broader moves to "deconstruct" multimodal approaches, generate evidence on specific therapeutics, understand mechanisms of action, and the growth in personalised medicine.

Psychoeducation/sleep hygiene

Psychoeducation: This typically comprises foundational information on the roles and functions of sleep, age-related changes, and how sleep–wake (circadian) patterns are regulated (e.g. the two-process model of sleep regulation; Borbély, 1982). Importantly, psychoeducation is usually integrated into elements of CBT-I rather than being standalone information. For example, the two-process model can be used to explain to patients that "bad nights" are often followed by "good nights" due to sleep homeostasis, serving to reinforce the rationale for sleep restriction and SCT.

Sleep hygiene: These recommendations (first presented by Hauri, 1991) are also reinforced in CBT-I, and comprise advice on bedroom factors (e.g. temperature, light and noise levels, bed comfort) and habits (e.g. caffeine, alcohol, establishing a regular routine, stop clock-watching) that can impact on sleep. It should be noted, however, that sleep hygiene on its own does not constitute an evidence-based treatment for insomnia, and sleep hygiene behaviours sometimes have a function of "safety behaviours" in patients with insomnia, heightening the perceived threat of not being able to sleep.

Recently the "five principles of good sleep health" have been proposed, and these may form a useful introduction to the cognitive-behavioural approach (Espie, 2022b).

Relaxation therapy

Many of the earliest insomnia studies included some form of relaxation intervention to facilitate de-arousal (Baglioni et al., 2022). This was consistent with the widespread adoption, in behavioural psychotherapy, of relaxation as a re-conditioning agent. Empirical research on insomnia suggests that abbreviated progressive muscle relaxation, autogenic training and imagery exercises are most frequently used for the treatment of insomnia. At present there is no evidence that one or the other method is superior, but the range of possibilities provides a multitude of treatment options.

Sleep-restriction therapy (SRT)

The behavioural model of insomnia put forth by Spielman et al. (1987) identified time in bed extension as important in the maintenance of insomnia. Extending time in bed (including napping), often as a compensatory response to poor sleep, can lead to increased sleep latencies, sleep fragmentation, and increased variability in the timing of the sleep–wake pattern. SRT aims to restrict sleep opportunity and harness increased sleep pressure to consolidate sleep and regularise its occurrence (Spielman et al., 1987). Once sleep is consolidated (indexed by sleep efficiency thresholds of > 85% or 90%), the goal is to extend time in bed, often on a weekly basis, to arrive at a sleep opportunity that delivers nightly sleep need, improved sleep continuity, and optimised daytime functioning. There is, however, variation in the configuration of SRT in relation to defining the initial time in bed prescription, minimum time in bed (e.g. 4.5, 5 or 6 hr), position of the sleep window, and weekly titration parameters (Kyle et al., 2015). SRT is hypothesised to work through restricting, regularising and reconditioning sleep opportunity, which drives a cascade in cognitive, behavioural and physiological pathways to improve sleep and daytime functioning (Maurer et al., 2018; Spielman et al., 1987). Evidence from randomised–controlled trials (RCTs) shows that acute implementation of SRT increases sleepiness proximal to bedtime, decreases pre-sleep

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 6 Meta-analyses on the efficacy of CBT-I in chronic insomnia/insomnia disorder without comorbidities

Author (year)	Population	No. studies/patients	Intervention	Study endpoints	Effects on study endpoints
Morin et al. (1994)	Insomnia	59/2102	CBT-I and single components	SOL, WASO, NOA, TST	(a) Good effects of CBT-I on all parameters (b) Good follow-up results
Murtagh and Greenwood (1995)	Insomnia	66/2007	CBT-I and single components	SOL, NOA, TST, SQ	(a) Good effects of CBT-I on all parameters (b) Good follow-up results
Pallesen et al. (1998)	Insomnia, age > 50 years	13/388	CBT-I and single components	SOL, NOA, WASO, TST	(a) Good effects of CBT-I on all parameters (b) Good follow-up results
Montgomery and Dennis (2004)	Primary insomnia, age > 60 years	7/322	CBT-I, bright light and physical exercise	SOL, TST, SE, WASO	(a) Good effects of CBT-I on sleep maintenance (b) Almost no effects of bright light and physical exercise
Irwin et al. (2006)	Insomnia, age > 55 years versus younger patients	23/NA	CBT-I and single components	SQ, SOL, TST, SE, WASO	Medium to strong effects in older patients
Okajima et al. (2011)	Primary insomnia	14/927	CBT-I	SOL, WASO, EMA, SE, PSG, ACT	(a) Good effects of CBT-I on all parameters (b) Good follow-up results
Miller et al. (2014)	Primary insomnia	4/192	SRT	SOL, WASO, TST, NOA, SE, SQ	Sleep restriction alone is effective
Koffel et al. (2015)	Insomnia	8/659	Group CBT-I	SOL, WASO, SE, SQ, TST, pain, depression	Group CBT-I is effective
Trauer et al. (2015)	Chronic insomnia	20/1162	CBT-I	SOL, WASO, TST, SE	Clinically relevant efficacy without undesired side-effects
Ballesio et al. (2018)	Insomnia	47/4317	CBT-I	Depression, fatigue	Small effects for depression, no significant effects for fatigue
Chung et al. (2018)	Insomnia	15/1194	Psychoeducation/ sleep hygiene versus CBT-I	ISI, SE, TST, SOL, WASO, AKT, PSQI	Psychoeducation/sleep hygiene less effective than CBT-I for PSQI, ISI, SE, SOL, WASO (medium to large ES)
Mitchell et al. (2019)	Insomnia disorder	15/1541	CBT-I	PSG, ACT	Small effects on actigraphy, no effects on PSG parameters
Van der Zweerde, van Straten, et al. (2019)	Insomnia	30/2835	CBT-I	SE, SOL, ISI 3, 6 and 12 months after therapy	Small to medium ES for ISI and SE, small ES for SOL at follow-ups
Benz et al. (2020)	Insomnia disorder	86/15,578	CBT-I	Daytime impairments in the context of insomnia	Small to medium ES for depression, anxiety, fatigue, quality of life and daytime functioning
Thakral et al. (2020)	Insomnia	16/1964	CBT-I	Dysfunctional thoughts and beliefs about sleep	Large ES for dysfunctional thoughts and beliefs about sleep post-treatment and at follow-up

(Continues)

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 6 (Continued)

Author (year)	Population	No. studies/patients	Intervention	Study endpoints	Effects on study endpoints
Ballesio et al. (2021)	Insomnia	15/1058	CBT-I	Repetitive negative thinking (e.g. ruminations)	Small ES for worry, no significant effect for rumination
Edinger et al. (2021)	Chronic insomnia	89/not indicated	CBT-I and its single components	TST, SE, SOL, WASO, SQ, ISI, PSQI	Medium to large ES for sleep parameters, smaller ES for single components
Kwon et al. (2021)	Insomnia > 60 years	28/2391	CBT-I, BT, acupuncture	PSQI	All treatments effective against waitlist
Maurer, Schneider, et al. (2021)	Insomnia	8/533	Sleep restriction	TST, SE, SOL, WASO, ISI	Large effects on SE, WASO, ISI; medium effects on SOL; no effect on TST
Xu et al. (2021)	Insomnia	31/2449	F2F CBT-I	TST, SE, SOL, WASO, NOA, ISI, PSQI, depression, anxiety, fatigue, somatic and mental health	Significant effects for ISI, PSQI, TST, SE, SOL, WASO, NOA, depression and fatigue; no significant effects on anxiety and mental health
Yu et al. (2021)	Insomnia	14/2263	CBT-I	TST, SE, SOL, depression, anxiety	Significant effects for TST, SE, SOL, depression and anxiety
Alimoradi et al. (2022)	Insomnia	24/1977	CBT-I	Quality of life	Small to medium effects on quality of life
Huang, Li, et al. (2022)	Insomnia > 60 years	14/792	CBT-I	TST, SE, SOL, WASO	Significant effects for TST, SE, SOL, WASO
Jansson-Fröjmark et al. (2022)	Insomnia	10/384	Paradoxical intention	TST, SE, SOL, NOA	Large ES for SOL, NOA, medium ES for TST, no effect for SE
Kwon et al. (2022)	Insomnia	10/496	Brief behavioural therapy	TST, SE, SOL, WASO	Significant effects for SE, SOL, WASO 1–8 weeks after therapy

Abbreviations: ACT, actigraphy; BT, behavior therapy; CBT-I, cognitive-behavioural therapy for insomnia; EMA, early-morning awakening; ES, effect size; F2F, face-to-face; ISI, Insomnia Severity Index; NOA, number of awakenings; PSG, polysomnography; PSQI, Pittsburgh Sleep Quality Index; SE, sleep efficiency; SOL, sleep-onset latency; SQ, sleep quality; SRT, sleep-restriction therapy; TST, total sleep time; WASO, wake time after sleep onset.

arousal, and improves markers of sleep continuity and sleep depth (Maurer, Schneider et al., 2022).

Stimulus control therapy (SCT)

The basic concept in stimulus control is that many patients with insomnia exhibit a learned association between the bed/bedroom and being awake (instead of sleeping). Bootzin (1972) conceived of SCT as an operant conditioning paradigm; we go to bed to get the reward of sleep. There is heavily reinforced instrumental learning in good sleepers because of the regular proximal relationship between bed and (rapid) sleep onset. However, this connection is broken in insomnia and counterproductive classical conditioning can occur. SCT comprises seven primary instructions with the aim of eliminating these maladaptive connections and reestablishing the connection where "bed" again equals "sleep" to the patient.

Cognitive therapeutics

Difficulties with cognitive arousal, mental events intruding upon sleep, heightened emotion and counterproductive sleep effort are very common in insomnia. Over the past few decades, numerous cognitive techniques have been developed and tested to address these problems. In most trials, and in routine clinical practice, these are commonly "bundled" within the "C" component of CBT-I. The interventions themselves are described in detail elsewhere (Baglioni et al., 2022; Espie, 2022a). In brief, they are cognitive control ("putting the day to rest" before going to bed), paradoxical intention (abandoning effort to sleep/attempting to remain awake), imagery training (active visualisation exercises that capture attention and manage rumination) and cognitive restructuring (challenging negative thoughts by means of Socratic dialogue or within the framework of behavioural experiments).

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

The evidence-base: CBT-I for insomnia without comorbidities

Meta-analyses concerning CBT-I are depicted in Table 6.

Up to now, 25 meta-analyses have been published focusing on the effect of CBT-I on insomnia/insomnia disorder in those without comorbidities since 1994. It should be critically mentioned that many early studies used a waitlist group as a control group, which is not as powerful as a placebo group, typically used in pharmacological studies. Furthermore, these meta-analyses are not independent of each other, thus there may be overlap within the studies. These issues, therefore, may overemphasise the effects of CBT-I. That said, a number of early placebo-controlled studies with small sample sizes (Espie et al., 1989; Turner & Ascher, 1979) exist, and suggested positive effects of CBT-I beyond sham comparison groups.

Most recent meta-analyses on the efficacy of CBT-I in insomnia/insomnia disorder, without comorbidities, show large effects on the severity of insomnia symptoms (see summary by Edinger et al., 2021), whereby follow-up measurements up to a year later show small to medium effects (van der Zweerde, Bisdounis, et al., 2019).

The evidence-base: CBT-I component-related efficacy

In relation to the monotherapy of insomnia, from the active elements of CBT-I, data up to now suggest efficacy of SRT (Edinger et al., 2021; Kyle et al., 2023; Maurer, Schneider, et al., 2021), SCT (Edinger et al., 2021), relaxation therapy (Edinger et al., 2021) and paradoxical intention as one specific cognitive therapeutic (Jansson-Fröjmark et al., 2022). Psychoeducation/sleep hygiene alone does not appear to be very effective (Chung et al., 2018; Edinger et al., 2021), probably because many patients are already aware of these recommendations (Lacks & Rotert, 1986). Thus, sleep hygiene advice given alone is not recommended in the treatment of chronic insomnia disorder and, in RCTs among patients with insomnia, sleep hygiene is often used as the "placebo" condition (Bjorvatn et al., 2011).

A network meta-analysis (Steinmetz et al., 2022, 2023) that aimed to delineate the efficacy of the different components of CBT-I showed that sleep restriction and stimulus control seem to be the most effective components of CBT-I. Nevertheless, interventions based on single strategies should be offered in full consideration of motivational and safety issues. Especially the behavioural strategies because sleep restriction and stimulus control may be very challenging for the patients; thus, the intervention may require careful consideration of the patient's motivation and readiness to apply the strategy with systematic regularity. Furthermore, these therapeutics may temporarily increase daytime sleepiness and fatigue in the early interventional phases, thus safety issues should be fully discussed with the patient.

The evidence-base: CBT-I effects beyond sleep complaints

Importantly, beyond beneficial effects on sleep-related outcomes, CBT-I also has positive effects on associated subclinical depressive symptoms, anxiety, daytime sleepiness and fatigue (Benz et al., 2020), dysfunctional cognitions related to sleep (Thakral et al., 2020), worry (Ballesio et al., 2021), and quality of life (Alimoradi et al., 2022).

The evidence-base: CBT-I for insomnia with comorbidities

The CBT-I is also effective when insomnia is comorbid with other medical disorders (Table 7).

There have been 15 published meta-analyses concerning CBT-I for insomnia when comorbid with other conditions since 2011.

In the area of mental disorders, medium to large effects have been reported on insomnia severity when insomnia is comorbid with depression, posttraumatic stress disorder or alcohol dependency (Hertenstein et al., 2022). There is not sufficient evidence concerning insomnia comorbid with bipolar disorders (Bisdounis et al., 2022) or psychotic disorders (Hertenstein et al., 2022). CBT-I is also effective when insomnia is comorbid with tinnitus (Curtis et al., 2021), chronic pain (Selvanathan et al., 2021), cancer (Gao, Liu, et al., 2022; Squires et al., 2022), sleep apnea (COMISA; Sweetman et al., 2023), and in patients with neurological disorders such as traumatic brain injury, stroke or Parkinson's disease (Ford et al., 2023; Lebrun et al., 2020). Interestingly, especially in the area of mental disorders, the evidence indicates that CBT-I not only has sleep-related effects, but also has a positive effect on the symptoms of the comorbid disorder/condition (Hertenstein et al., 2022).

The evidence-base: CBT-I in self-help formats or delivered digitally

Over the past 10 years there has been a growing literature on the application of CBT-I in self-help formats, particularly using either self-help books/texts or, now, dCBT-I. To date, 11 meta-analyses are available on this topic (Table 8). A meta-analysis by Gao, Ge, et al. (2022), explored a range of delivery formats, and reported that CBT-I in single therapy, in group therapy and in digital form, with or without support by therapists, is effective. Single therapy, group therapy and digital therapy with personal support were most effective (Gao, Ge, et al., 2022). A further recent meta-analysis by Hasan et al. (2022) also suggests that personal support is to be preferred when CBT-I is administered digitally.

Interestingly, the most contemporary network meta-analysis on all forms of CBT-I administrations (i.e. onsite versus digital and other settings; Simon, Steinmetz, et al., 2023) demonstrated that individual F2F CBT-I, group F2F CBT-I, telehealth (videoconference, phone call) and guided bibliotherapy conveyed the strongest effects on insomnia, whereas guided and unguided dCBT-I yielded medium effect sizes, slightly favouring guided over unguided dCBT-I. Smart phone applied CBT-I did not attain significant effects compared with control.

In Europe, several dCBT-I treatment programs are now available in web and/or mobile application format. Table 9 presents information on these programs, their evidence base, and levels of adoption.

To obtain this information, all members of the EIN ($n = 250$) and all authors of this guideline ($n = 44$) were asked to complete a questionnaire about digital tools available in their home country, which resulted in the above-mentioned tables. The evidence base reported reflects publications available from peer-review journals. For a systematic overview encompassing the wider number of apps claiming to improve sleep that are available through App stores, see Simon, Reimann, et al. (2023).

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 7 Meta-analyses on the efficacy of CBT-I on insomnia/insomnia disorder with comorbidities

Author (year)	Population	No. studies/patients	Intervention	Study endpoints	Effects on study endpoints
Belleville et al. (2011)	Insomnia with/without comorbid anxiety	50/2690	CBT-I	Anxiety scales	Moderate effects on anxiety
Geiger-Brown et al. (2015)	Comorbid insomnia (somatic/mental)	23/1379	CBT-I	SOL, WASO, TST, SE, ISI, PSQI	Good efficacy; long-term effects at 18 months
Wu et al. (2015)	Comorbid insomnia (somatic/mental)	37/2189	CBT-I	SOL, WASO, SQ, TST, remission, comorbid symptoms	Good efficacy; smaller effects on comorbid symptoms; better effects for mental outcomes
Ho et al. (2016)	Insomnia + PTSD	11/593	CBT-I	SOL, WASO, SE, TST, PTSD symptoms	Good sleep effects, good effects on PTSD symptoms
Johnson et al. (2016)	Insomnia + cancer	8/752	CBT-I	SE, WASO, ISI, cancer symptoms	Good sleep effects, good effects on cancer symptoms
Tang et al. (2015)	Insomnia + pain	11/1066	CBT-I	SQ, fatigue, pain	Good sleep effects, good effects on comorbid symptoms
Van Straten et al. (2018)	Insomnia, with and without comorbidities	87/6303	CBT-I	SE, SOL, ISI	Significant medium to large ES for SE, SOL, ISI
Feng et al. (2020)	Insomnia disorder + depression	17/1756	CBT-I	ISI, PSQI, depressive symptoms	Significant ES for ISI/PSQI, weaker effects for depressive symptoms
Zhou et al., 2020	Insomnia disorder + psychiatric/somatic comorbidity	13/853	CBT-I	SOL, WASO, SQ, ISI	Medium ES for ISI, SQ; weaker effects for SOL, WASO
Curtis et al. (2021)	Insomnia + tinnitus	4/470	CBT-I	ISI	Medium ES for ISI
Ma et al. (2021)	Insomnia disorder + breast cancer	14/1363	CBT-I	ISI	Medium ES for ISI
Selvanathan et al. (2021)	Insomnia + chronic pain	12/762	CBT-I	Global sleep parameters, pain, depressive symptoms	Strong ES for global sleep parameters, smaller ES for pain and depressive symptoms
Gao, Liu, et al. (2022)	Insomnia + cancer	16/1523	CBT-I	SE, SOL, WASO, ISI	Small to medium ES for ISI, SOL, WASO
Hertenstein et al. (2022)	Insomnia + comorbid mental disorders	22/1083	CBT-I	ISI, comorbid symptom severity	Significant medium to large ES for ISI, significant ES for symptoms of comorbid disorder
Squires et al. (2022)	Insomnia + cancer	22/1461	CBT-I	ISI, TST, SE, SOL, WASO, SQ, anxiety, depressive symptoms, fatigue, quality of life	Medium ES for ISI, small to large ES for TST, SE, SOL, WASO, SQ, anxiety, depressive symptoms, fatigue, quality of life

Abbreviations: CBT-I, cognitive-behavioural therapy for insomnia; ES, effect size; ISI, Insomnia Severity Index; PSQI, Pittsburgh Sleep Quality Index; PTSD, posttraumatic stress disorder; SE, sleep efficiency; SOL, sleep-onset latency; SQ, sleep quality; TST, total sleep time; WASO, wake time after sleep onset.

Table 9 summarises dCBT-I treatments where effectiveness for insomnia is supported through at least one pre-registered published randomised–controlled trial (RCT). As described in the table, some programs incorporate guidance from health professionals whilst others are fully automated. Guidance means participants are offered support (either by mail or phone) to discuss critical issues and progress. Almost half of the evidence-based dCBT-I programs are now reimbursed by the local/national health authorities, and can be recommended or prescribed at the cost of local health insurance. Sleepio from the UK (also available in the USA) is the most extensively researched dCBT-I application, having generated 13 published RCTs including a placebo-controlled trial.

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 8 Meta-analyses on the efficacy of self-help/digital CBT-I for insomnia

Author (year)	Population	No. studies/patients	Intervention	Study endpoints	Effects on study endpoints
Van Straten and Cuijpers (2009)	Insomnia	10/1000	Self-help CBT-I	SOL, WASO, SE, SQ, TST	Small to moderate effects
Cheng and Dizon (2012)	Insomnia	6/433	dCBT-I	SOL, WASO, SE, SQ, TST	Small to moderate effects
Ho et al. (2015)	Insomnia	20/2411	Self-help + dCBT-I	SOL, WASO, SE, SQ, TST	Self-help CBT-I is effective and acceptable as a starter for treatment
Ye et al. (2015)	Insomnia with comorbid conditions	9/776	dCBT-I	Anxiety, depression	Moderate effect sizes for comorbid symptoms
Zachariae et al. (2016)	Insomnia	11/1460	dCBT-I	ISI, SOL, WASO, NOA, TST, SQ	Comparable to F2F CBT-I
Seyffert et al. (2016)	Insomnia	15/2392	dCBT-I	ISI, SOL, TST, WASO, NOA, SQ, PSQI	Good efficacy for sleep parameters, good follow-up results
Soh et al. (2020)	Insomnia	33/9364	dCBT-I	ISI, other sleep parameters	Small ES for ISI
Ho et al. (2020)	Insomnia + depression	30/5945	Self-help CBT-I	Insomnia symptoms, depressive symptoms	Medium ES for insomnia symptoms, smaller ES for depressive symptoms
Gao, Ge, et al. (2022)	Insomnia	61/11,571	CBT-I in several formats including dCBT-I	TST, SE, SOL, WASO, ISI	Significant effects of individual and group CBT-I and dCBT-I with or without assistance; largest for individual and group CBT-I and dCBT-I with assistance
Hasan et al. (2022)	Insomnia	54/11,815	dCBT-I	TST, SE, SOL, WASO	Significant effects of assisted dCBT-I for TST, SE, SOL, WASO
Forma et al. (2022)	Chronic insomnia	20/5659	dCBT-I	SOL, WASO, ISI	Significant effects for ISI, WASO

Abbreviations: CBT-I, cognitive-behavioural therapy for insomnia; dCBT-I, digital-cognitive behavioural therapy for insomnia; ES, effect size; F2F, face-to-face; ISI, Insomnia Severity Index; NOA, number of awakenings; PSQI, Pittsburgh Sleep Quality Index; SE, sleep efficiency; SOL, sleep-onset latency; SQ, sleep quality; TST, total sleep time; WASO, wake time after sleep onset.

Table S1 (supplemental material) summarises programs which, up to now, lack evidence through an RCT. Some of these programs have also been tested in uncontrolled single-arm trials.

To summarise, at present there is a rapidly developing market of digital health applications targeting the treatment of insomnia, with some of the applications already having a considerable database and being reimbursed by local health authorities. Concerning RCT supported applications (Table 9), five European languages (including Dutch, English, French, German and Swedish) are already covered. This development definitely reflects the overall trend for digitalisation in healthcare, but may also be seen as a consequence of our previous guideline (Riemann, Baglioni, et al., 2017; Riemann, Baum, et al., 2017), in which we recommended CBT-I as first-line treatment for insomnia. Digital CBT-I surely will make access to CBT-I easier than before (particularly when fully automated and 24/7 available), meeting the needs of many patients with insomnia who have no access to CBT-I therapists. In any case, at present, it must be stressed that this type of therapy has to undergo rigorous quality control as well (e.g. does it really contain all important ingredients of CBT-I?) and is it evidence-based, meaning has it been subjected to at least one high-quality RCT? Gold-standard evidence criteria for the regulation of dCBT-I (digital therapeutics) have been published (Espie, Torous, & Brennan, 2022; Table 10). Importantly, developers should be aware that an evidence-informed intervention (broadly based on CBT-I but not specifically subjected to trials) is not an adequate substitute for being evidence based (Espie, Firth, & Torous, 2022).

We also stress that local health authorities in the various European countries should take over this regulatory function—in Germany and the UK, for example, the same authority that regulates medications has embraced this task. Future research will have to demonstrate whether digital CBT-I is really equivalent to in-person/F2F therapy and to which degree increased rates of dropouts/attrition may occur, especially in cases of comorbid insomnia. Caution is urged in cases of comorbid insomnia, especially when severe

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 9 dCBT-I applications with scientific evidence (at least one RCT) in Europe

Language (country of origin)	Name of product	Web address	Computer/app based	Reimbursement by national healthcare	Reference (s)
Dutch The Netherlands	i-Cycle	https://slaapregister.nl	Yes/Yes (partly) + assistance[a]	No	Leerssen et al. (2022)
Dutch The Netherlands	Somnio	https://somnio.nl	Yes/Yes + assistance[a]	Yes	Dekker et al. (2020)
Dutch The Netherlands	i-Sleep	https://www.minddistrict.com https://therapieland.nl https://leerslapen.nl	+ assistance[a]	Yes	Baka et al. (2022); Van der Zweerde, van Straten, et al. (2019); Van der Zweerde et al. (2020); Van Straten et al. (2014)
English UK	SLEEPIO	https://www.sleepio.com/	Yes/Yes	Yes (Scotland)	More than 30 publications; Selection: Espie et al. (2012, 2019); Espie, Kyle, Miller, et al. (2014); Felder et al. (2020); Freeman et al. (2017)
French France	TheraSomnia	https://www.therasomnia.com/	Yes/Yes + assistance[a]	No	Lopez et al. (2019)
German Austria	NUKKUAA	www.nukkuaa.com	No/Yes	No	Schabus et al. (2022); Eigl et al. (2022); Hinterberger et al. (2022, 2023)
German Germany	SOMNIO	https://somn.io/	Yes/Yes	Yes	Lorenz et al. (2019)
German Germany	HelloBetter Schlafen/ HelloBetter sleep	https://hellobetter.de/online-kurse/schlafen/	+ assistance[a]	Yes	Behrendt et al. (2020); Ebert et al. (2015); Thiart et al. (2015)
German Switzerland	Meinstresscoach/ SweetDreams	https://www.meinstresscoach.ch/kurse	No/Yes + assistance[a]	No	Hürlimann et al. (2023)
Swedish Sweden	Internetpsykiatri Sömnproblem—insomni	https://www.internetpsykiatri.se/behandling/somnproblem-insomni/	Yes/No + assistance	Yes	Kaldo et al. (2015); Blom et al. (2016)
Swedish Sweden	Livanda Sömnproblem (insomnia)	https://www.livanda.se/kbt-internetterapi-somnproblem.aspx	Yes/No + assistance[a]	No	Ström et al. (2004)

[a]Assistance: mail/phone interaction with a health professional possible.

acute mental disorders are present. It is important that the pathways to the various CBT-I formats are based on professional expertise. Also, the question of adverse events and how these are dealt with needs to be considered thoroughly (in all CBT-I programs). Finally, sleep trackers, with often inaccurate sleep feedback, need to be more thoroughly tested and should in general be discouraged for patients with insomnia until published validation of their accuracy against PSG is available.

Side-effects and contraindications of CBT-I

Typical of the psychotherapy literature more generally, adverse effects of CBT-I have received limited attention in clinical trials (Condon et al., 2021). This situation, however, is changing as trial methodology and reporting improve. Clinical experience shows that SRT is coupled with increased tiredness and sleepiness, and there is evidence from both controlled and uncontrolled trials of objectively reduced vigilance during the first weeks of implementation (Kyle et al., 2014; Maurer et al., 2022; Maurer, Schneider, et al., 2021). Indeed, the initial restriction of time in bed and its associated sleepiness may be central to therapeutic effects of behavioural treatment (Maurer et al., 2020). In a large trial ($n = 1711$) of dCBT-I, Espie and colleagues (2019) showed that CBT-I was associated with higher reports of intervention-attributed adverse effects (extreme sleepiness, fatigue or exhaustion, attention and concentration problems, low motivation and energy, headaches, memory difficulties and irritability) relative to sleep hygiene. Further work by Sunnhed et al. (2020) localised adverse effects, especially fatigue and sleepiness, to behavioural therapy (SRT, SCT) relative to cognitive therapy. As such, care should

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 10 WHAT criteria need to be met by a dCBT/digital therapeutic program? (from Espie, Torous & Brennan 2022)

dCBT criteria	Demonstrated by …
W: The therapeutic has to WORK	Treatment efficacy needs to be supported by adequately powered RCT(s) to demonstrate level 1 evidence of change that is attributable to the digital therapeutic. The trial and its primary and secondary outcomes should be pre-registered, and those data should be published following peer-review
H: The therapeutic has to HELP	Statistically significant between-group effects of the digital therapeutic over a control condition are important. However, RCT(s) need to demonstrate clinically meaningful differences or clinical remission endpoints relating to the target therapeutic condition as assessed by clinically accepted and standardised pre-registered measures
A: The therapeutic has to be ACCESSIBLE	A digital therapeutic has to use high-quality technology to deliver a reliable and acceptable patient experience within the target population. There needs to be evidence that the digital therapeutic is capable of scaling safely and effectively to population level distribution in the real world
T: The therapeutic has to be TRUSTED	The digital therapeutic needs to adhere to clinical governance and risk management standards for software as a medical device, privacy and data security certifications, and reflect national and international clinical guidelines for information storage and data transfer

Abbreviations: dCBT, digital cognitive-behavioural therapy; RCT, randomised–controlled trial.

be taken when administering SRT and/or stimulus control protocols, including initial assessment for suitability and ongoing reviews of sleepiness (and impact on driving) during treatment, with appropriate adaptations put in place where necessary. In the largest test of SRT to date ($n = 642$), Kyle et al. (2023) found no evidence that nurse-delivered SRT was associated with serious adverse events, or increased incidence of important pre-defined adverse events (falls, work-related or motor vehicle accidents, near-miss driving incidents, or falling asleep while driving) relative to sleep hygiene control. Contraindications for treatments like stimulus control or sleep restriction that may entail partial sleep deprivation include any kind of epilepsy and conditions that might be aggravated by sleep loss. It should be noted of course, that as with medications, patients and clinicians should generally be familiar with risk–benefit considerations and associated decision-making.

Up to now, no mention has ever been made that CBT-I is associated with any risks of abuse/dependency, "dose increase", tolerance or rebound insomnia—all problems associated with some of the hypnotic medications. The low probability of serious or enduring adverse effects with CBT-I is considered a strong advantage over medications.

Combining CBT-I with pharmacotherapy

Concerning the question of whether a combination of CBT-I and pharmaceutical treatment is clinically superior to either therapy in isolation, there are two high-quality original studies, from some years ago, using temazepam or zolpidem as medications (Morin et al., 1999, 2009). Both studies showed that during the acute treatment phase a synergistic effect of CBT-I and pharmacotherapy can be observed; however, after acute therapy mono-treatment with CBT-I is superior to the combination of CBT-I and pharmacotherapy. More recent work demonstrated that sequential treatment, starting with behavioural therapy and subsequently zolpidem, resulted in the largest percentage response and remission in the longer term (1 year), indicating that secondary administration of medication could be an effective strategy in insomnia management (Morin et al., 2020). This high-quality controlled study also found that approximately half of the patients remitted with these interventions, underscoring the urgency for the development of more effective treatments, as opposed to CBT-I or pharmacotherapy, for the other half. This importance is further underlined by one of the few meta-analyses that reports actual post-CBT-I sleep efficiency scores, as opposed to effect sizes only (Seyffert et al., 2016): sleep efficiency does not normalise in more than half of those treated.

Insomnia care in Europe: the role of CBT-I

In conclusion for this section, the published evidence suggests that CBT-I should be used as first-line treatment for patients with chronic insomnia. This recommendation is independent of the question whether the insomnia is the sole presenting problem, or comorbid to another medical disorder. Single components (especially sleep restriction and stimulus control) may be effective on their own, and there is evidence that brief treatments (consisting of one–two sessions) may also have some effect on their own (Kwon et al., 2022). A major advantage of CBT-I over pharmacological treatment is its treatment sustainability—in contrast to pharmacotherapy where symptoms frequently recur or even rebound after treatment discontinuation, where no such effect has been reported for CBT-I. Furthermore, as suggested by the evidence, the effects of CBT-I (Tables 6–8) appear to be stable over time (follow-ups up to 10 years have been published; Jernelöv et al., 2022) and may only gradually weaken. Booster sessions may be needed to refresh the initial treatment gains.

However, a huge problem presently in healthcare across Europe is the fact that up to now in-person/F2F CBT-I is not available for all patients suffering from insomnia. It needs to be seen whether this gap in clinical care can be closed by digitally based interventions. The accelerating and substantial evidence from trials of dCBT-I applications are very promising and may greatly improve guideline-based insomnia care in Europe.

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Another important issue at this point (which also applies to pharmacotherapy) is the question as to what extent CBT-I (in all its forms of applications) is associated with attrition/adherence, and at what point are patients considered mere responders (showing some meaningful clinical benefit) or remitters (being fully resolved, i.e. not fulfilling threshold values, e.g. on the ISI). These issues are especially pertinent for fully automated digital treatments, i.e. applied without any human guidance, where reported dropout rates have been high (> 50%). It is recommended that future RCTs and, even more so, comparative meta-analyses correct their effect sizes for the differential attrition they have (i.e. intention to treat). Few studies have investigated the predictors of dropout. As such, it is imperative to determine who are most likely to drop out of F2F or dCBT-I, and which alternative treatments they might best respond to.

Future research will also have to address the issue of head-to-head comparisons between all types of treatments (both psychological and/or pharmacological). Furthermore, beyond well-controlled randomised clinical trials we are in need of data from real-world everyday clinical practice regarding treatment acceptance, adherence, attrition, side-effects and outcome.

4.1.2 | Other cognitive-behavioural/psychotherapeutic interventions

As previously mentioned, the range of treatments for insomnia has expanded over the past few decades, and these have often been combined into the multicomponent approach. Recent expansions in this literature have included the so-called "third-wave" therapies such as mindfulness, and acceptance and commitment therapy (ACT; n.b. "first-wave": behavioural; "second-wave": cognitive).

Two contemporary meta-analyses concerning mindfulness-based treatments for insomnia show unequivocally positive effects (Chen, Chang, et al., 2020; Wang, Wang, et al., 2020). The number of studies in this area, however, is still relatively small, and the authors of both meta-analyses stress that long-term effects need to be investigated more thoroughly. ACT has been studied only in small randomised clinical studies. While ACT was shown to be superior to an active control condition (Zakiei et al., 2021), it was inferior to CBT-I (El-Rafihi-Ferreira et al., 2022). Moreover, replacing the cognitive therapy aspect of CBT-I with ACT did not enhance efficacy (El Rafihi-Ferreira et al., 2021). Finally, one study of a web-based treatment platform using ACT, SRT and SCT found it was superior to waitlist control conditions (Chapoutot et al., 2021). In summary, there is evidence that ACT may be effective in insomnia; however, larger randomised clinical studies are desperately needed.

A meta-analysis by Lam et al. (2015) on hypnotherapy for insomnia suggested some positive effects. The methodological quality of the original studies, however, was judged as poor, which casts doubt on the benefits of this approach at this time.

Intensive sleep retraining is a newer form of behavioural treatment, requiring PSG monitoring. Scott et al. (2023) summarise promising evidence for this new type of training; however, at present data are still too scarce to provide a recommendation.

Summarising, further research is needed for these other cognitive-behavioural forms of psychotherapy before they can be recommended for the treatment of insomnia.

4.1.3 | Exercise, light therapy, music and non-invasive brain stimulation

Concerning the efficacy of exercise in the treatment of insomnia, there is some evidence; however, further high-quality studies are necessary (published meta-analyses: Amiri et al., 2021; Baglioni, Bostanova, et al., 2020; Banno et al., 2018; Choong et al., 2022; D'Aurea et al., 2022; Li, Li, et al., 2021; Mercier et al., 2017; Rubio-Arias et al., 2017 T). One RCT showed that the addition of exercise to CBT-I helped maintain its initial beneficial effects and this was largely due to CBT-I (Dekker et al., 2020). Tai-Chi is a specific form of exercise suggested to be effective by most recent meta-analyses; however, it was highlighted that more high-quality studies are still needed (Han et al., 2022; Li, Chen, et al., 2020).

A meta-analysis on light therapy for insomnia showed small positive effects on sleep (Van Maanen et al., 2016) and this is an area worthy of further investigation, especially because a circadian sleep–wake rhythm disorder is frequently comorbid with insomnia. Chambe et al. (2023) came to similar conclusions: light therapy may have some promise, but further RCTs are necessary. One RCT showed that the addition of light to CBT-I helped maintain its initial beneficial effects due to CBT-I (Dekker et al., 2020).

The most recent meta-analyses on the effects of listening to music on sleep in patients with insomnia also suggest a possible positive effect (Chen et al., 2021; Feng et al., 2018; Jespersen et al., 2015). However, there are few original studies on these topics and the quality of the available literature was criticised in these meta-analyses.

Several non-invasive brain stimulation techniques have been tested as potential treatment approaches for insomnia, in particular repetitive transcranial magnetic stimulation (rTMS), transcranial electric stimulation (tES) comprising transcranial direct current stimulation and transcranial alternating current stimulation, as well as vagus nerve stimulation and brain cooling. While several RCTs report positive results, methodological issues such as the lack of blinding or insufficient sham controls limit their interpretability. The therapeutic benefits of current non-invasive brain stimulation protocols for insomnia treatment are likely overestimated and, at present, no recommendation can be given for the use of brain stimulation approaches as a treatment strategy for insomnia (Krone et al., 2023).

In light of the available evidence, or lack thereof in some cases, it is considered premature to recommend any of these treatments as standalone interventions for insomnia. We do, however, suggest that elements of exercise and light therapy be integrated with CBT-I and may deliver additional benefits.

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 11 Major drug classes used to treat insomnia in Europe

BZ	Diazepam, flunitrazepam, flurazepam, lormetazepam, nitrazepam, oxazepam, temazepam, triazolam
BZRA	Zaleplone, zolpidem, zopiclone, eszopiclone
Sedating antidepressants	Agomelatine, amitriptyline, doxepin, mianserin, mirtazapine, trazodone, trimipramine
Antipsychotics	Chlorprothixene, levomepromazine, melperone, olanzapine, pipamperone, prothipendyl, quetiapine
Antihistamines	Diphenhydramine, doxylamine, hydroxyzine, promethazine
Phytotherapeutics	Hops, kava-kava, melissa, passiflora, valerian, lavender
Melatonin receptor agonists	Fast-release melatonin, ramelteon, PR melatonin
Orexin receptor antagonists	Daridorexant

Abbreviations: BZ, benzodiazepines; BZRA, benzodiazepine receptor agonists; PR, prolonged-release.

4.2 | Pharmacological treatments

A summary of the available and often used medications (on-label and off-label) for insomnia in Europe is provided in Table 11.

4.2.1 | Benzodiazepines (BZ) and benzodiazepine receptor agonists (BZRA)

The BZ and BZRA medications that are available in Europe are listed in Table 11. Table 12 summarises the meta-analyses on the efficacy of BZs and BZRAs.

Most of the meta-analyses demonstrate that BZ and BZRA drugs have positive effects on sleep when taken up to a maximum of 4 weeks. Beyond that timeframe, the meta-analyses by Liang et al. (2019) and Rösner et al. (2018) suggest positive effects of eszopiclone for a period of application of up to 6 months in certain cases, but this is to be decided on an individual basis.

Side-effects associated with BZs and BZRAs used include the development of tolerance and dependence, nocturnal confusion and falls (Treves et al., 2018), negative effects on cognitive functioning including memory impairment (Barker et al., 2004; Stranks & Crowe, 2014), hangover effects with associated impairments in driving capability, especially from substances with long half-lives (Rapoport et al., 2009), as well as rebound insomnia after withdrawal. Concerning the development of dependence from hypnotics, some authors postulate that in Germany up to a million people may be affected (Soyka, 2021).

Considering the risks of long-term use of these substances especially when considering the comparative efficacy of other treatment options, the scientific evidence, according to our evaluation, does not support a general positive recommendation for BZ and BZRA in the longer-term treatment of insomnia beyond 4 weeks. Although eszopiclone has been investigated in studies for 3 and 6 months and with long-term data for up to 12 months (see overview by Rösner et al., 2018), it should be used with caution when used beyond 4 weeks given its mechanism of action, as with other BZRAs. In general, longer-term treatment (mostly off-label use) with BZ or BZRA, either daily or preferably intermittently, may be initiated in some cases but the advantages and disadvantages need to be discussed on a case-by-case basis.

4.2.2 | Low-dose sedating antidepressants

The sedating antidepressants, which are frequently used in Europe to treat not only comorbid insomnia but also insomnia without comorbidities, are listed in Table 11. None of these substances has a general indication on the market for insomnia, in the absence of comorbid depression. Off-label use, therefore, is usually standard. Typically, sedative antidepressants are prescribed in insomnia at lower doses than are used in depression treatment. The meta-analytic literature on sedating antidepressants for insomnia is summarised in Table 13.

The meta-analysis by Everitt et al. (2018) suggests that there is only scant evidence for indicating sedating antidepressants in the treatment of insomnia. Significant but small effects were noted for doxepin and trazodone in the short term, up to 4 weeks. Undesired side-effects must always be considered carefully. Longer-term treatment of insomnia disorder (without comorbidities; off-label use) with low-dose sedating antidepressants may be initiated in some cases but the advantages and disadvantages need to be discussed on a case-by-case basis.

4.2.3 | Antipsychotics

The antipsychotic medications that are used quite frequently for the treatment of insomnia, especially where there are comorbid mental disorders, are listed in Table 11. The antipsychotics that explicitly mention insomnia, as an indication in the drug information, are melperone and pipamperone. There are, however, no randomised controlled clinical studies on these substances concerning insomnia disorder, either with or without comorbidities. Therefore, at present the scientific evidence does not recommend the use of antipsychotics (including quetiapine) in the treatment of insomnia without comorbidities, in either the short or long term.

4.2.4 | Dual orexin receptor antagonists (DORAs)

The introduction of DORAs has probably been the most significant recent development in the pharmacological treatment of insomnia. Among this family of drugs, the only one approved by the European

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 12 Meta-analyses on the efficacy of BZ and BZRA in the treatment of insomnia

Author (year)	Population	No. studies/patients	Intervention	Study endpoints	Effects on study endpoints
Nowell et al. (1997)	Primary insomnia	22/1894	BZ + zolpidem versus placebo, short-term treatment	SOL, NOA, TST, SQ	Significant improvement of sleep
Soldatos et al. (1999)	Insomnia	75/1276	Rapidly eliminated BZ + BZRA hypnotics versus placebo	SOL, TST, WASO (all PSG)	(a) Strong evidence for significant initial efficacy (b) Increased risk for development of tolerance and rebound phenomena; difference between substances
Holbrook et al. (2000)	Primary insomnia	45/2672	BZ + zopiclone versus placebo, short-term treatment	SOL, TST, USE	(a) Significant improvement of sleep (b) Increased risk for USE
Dündar et al. (2004)	Insomnia	24/3909	BZ versus BZRA, short-term treatment	SOL, TST, NOA, WASO, SQ, USE	(a) No difference between substances
Glass et al. (2005)	Insomnia, age > 60 years	24/2417	BZ + BZRA versus placebo, short-term treatment	SQ, SOL, TST, NOA, USE	(a) Significant improvement of sleep (b) Increased risk for USE
Buscemi et al. (2007)	Chronic insomnia	105/5582	BZ + BZRA + sedating antidepressants	SOL + secondary outcomes, USE	BZ and BZRA are effective; more USE with active drugs versus placebo
Huedo-Medina et al. (2012)	Insomnia	13/4378	BZRA (zolpidem, zaleplone, eszopiclone)	SOL + secondary outcomes	Small but significant effects on subjective and objective SOL
Winkler et al. (2014)	Insomnia	31/3820	BZ, BZRA, sedating antidepressants, melatonin	PSG and subjective sleep parameters	BZ and BZRA have significant effects on subjective and objective outcomes; smaller effects for antidepressants
Rösner et al. (2018)	Insomnia disorder	14/4732	Eszopiclone versus placebo	TST, SE, SOL, WASO	Significant effects for TST, SE, SOL, WASO, persisting up to 6 months
Liang et al. (2019)	Primary insomnia	6/2809	Eszopiclone versus placebo	TST, SOL, WASO, NOA	Significant effects for TST, SOL, WASO, NOA, persisting up to 6 months
Xiang et al. (2021)	Insomnia disorder	6/1068	Zolpidem versus placebo (treatment at least 1 month)	TST, SOL, WASO, SQ	Significant effects for TST, SOL, SQ, but not for WASO

Abbreviations: BZ, benzodiazepines; BZRA, benzodiazepine receptor agonists; NOA, number of awakenings; PSG, polysomnography; SE, sleep efficiency; SOL, sleep-onset latency; SQ, sleep quality; TST, total sleep time; USE, undesired side-effects; WASO, wake time after sleep onset.

Medicines Agency (EMA, 2022, 2023) so far has been daridorexant, which is indicated for treating adults with insomnia for a duration of least 3 months and has a considerable impact on how they function during the day (https://www.ema.europa.eu/en/medicines/human/EPAR/quviviq). This approval is mainly based on two controlled phase-III studies that compared daridorexant with placebo in 1854 patients with insomnia over a period of 3 months, and constitutes, so far, the largest clinical program among the available drugs in Europe. Over that period, a dose of 50 mg daridorexant showed significant effects on PSG-defined sleep parameters (sleep latency and wake after sleep onset) as well as increases in subjective total sleep time and reductions in daytime sleepiness/fatigue, with small to medium effect sizes (Mignot et al., 2022). While daridorexant improved objective (PSG-defined) parameters, no subjectively reported improvements in sleep-onset latency or sleep maintenance disturbances were observed. Overall, adverse effects were mild, and their incidence was comparable between daridorexant and placebo (with nasopharingitis, headache and somnolence being most common). Furthermore, an open extension study for up to 1 year in 804 patients demonstrated a maintained efficacy over this period, with no next-morning sleepiness and no withdrawal-related symptoms or rebound being observed after treatment discontinuation. Most importantly, no new safety problems, to the ones already known, arose during this long-term treatment period (Kunz et al., 2023).

These interesting data remain to be validated through practical experience in everyday practice. Considering that at the time of writing daridorexant has not yet been launched in most European countries, caution should prevail. In general, orexin receptor antagonists

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 13 Meta-analyses on the efficacy of sedating antidepressants in the treatment of insomnia

Author (year)	Population	No. studies/patients	Intervention	Study endpoints	Effects on study endpoints
Buscemi et al. (2007)	Chronic insomnia	105/873	BZ + BZRA + sedating antidepressants	SOL	Sedating antidepressants are less effective than BZ/BZRA
Winkler et al. (2014)	Insomnia	31/3820	BZ + BZRA + sedating antidepressants + melatonin	Subjective and objective sleep parameters	Sedating antidepressants are less effective than BZ/BZRA
McCleery et al. (2014)	Insomnia comorbid with M. Alzheimer	5/313	Trazodone + melatonin + ramelteon	SOL, TST, WASO, SE	Trazodone improves TST and SE
Yeung et al. (2015)	Insomnia	9/1983	Low-dose doxepin	Subjective and objective sleep parameters	Small to moderate effects for sleep maintenance and TST, but no effects for SOL
Liu et al. (2017)	Insomnia disorder	9/968	Tricyclic antidepressants	TST, SE, SOL, WASO (all PSG)	Medium effects for TST, WASO; smaller effects for SE, SOL; more USE than placebo
Everitt et al. (2018)	Insomnia	23/2806	Antidepressants	TST, SE, SOL (PSG), SQ	Tricyclics have significant small effects on SQ and on TST and SE, but not on SOL. Small effects of trazodone on various parameters; Side-effects have not been systematically studied.
Yi et al. (2018)	Insomnia disorder	7/429	Trazodone	PSG: SE, TST, SOL, WASO, NOA, SQ	Significant effects for NOA and SQ; none for SE, TST, SOL, WASO
Zheng et al. (2022)	Insomnia disorder	11/466	Trazodone	PSG: TST, SOL, WASO, NOA, N1, N2, SWS, REM	Significant increase of TST, SWS; significant reduction of SOL, WASO, NOA, N1; no effects on N2, REM

Abbreviations: BZ, benzodiazepines; BZRA, benzodiazepine receptor agonists; NOA, number of awakenings; PSG, polysomnography; REM, rapid eye movement; SE, sleep efficiency; SOL, sleep-onset latency; SQ, sleep quality; SWS, slow-wave sleep; TST, total sleep time; USE, undesired side-effects; WASO, wake time after sleep onset.

can be used for periods of up to 3 months. In individual cases, treatment can be extended for longer-term periods (up to 1 year) following discussion with the patient regarding the advantages and disadvantages reported in the 1-year study.

4.2.5 | Antihistaminergic drugs

Available antihistamines in Europe are depicted in Table 11. The table summarises over-the-counter (OTC) medication (diphenhydramine, doxylamin), and substances like hydroxyzine and promethazine that have to be prescribed by a physician. The status of these drugs (OTC versus prescription medicine) may vary from one European country to the other.

The evidence concerning antihistaminics in the treatment of insomnia is insufficient. One systematic review stated that there were no high-quality randomised–controlled clinical studies demonstrating efficacy compared with placebo—concluding that the evidence for these drugs is small and there might also be a rapid development of tolerance (Vande Griend & Anderson, 2012). Therefore, the scientific evidence at this point does not support a recommendation for antihistaminergic drugs in the treatment of insomnia either short or long term. Recently published evidence (Oyekan et al., 2021), indicating an association of antihistamines with increased mortality, suggests further caution.

4.2.6 | Melatonin

Melatonin is freely available in most European countries and can also be prescribed by physicians in most European countries for patients with insomnia. One substance (prolonged-release melatonin = PR melatonin) has been approved for the treatment of insomnia in patients older than 55 years (EMA, 2007/2010/2023—Circadin, https://www.ema.europa.eu/en/medicines/human/EPAR/Circadin). Ramelteon, a synthetic melatonin receptor agonist, is not available in Europe presently. The meta-analytic literature on melatonin in insomnia is shown in Table 14.

The meta-analysis by Marupuru et al. (2022) summarises studies on melatonin (including PR melatonin) and ramelteon. The authors report small to medium effects on sleep-related parameters in elderly

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 14 Meta-analyses on the efficacy of melatonin and melatonin receptor agonists in the treatment of insomnia

Author (year)	Population	No. studies/patients	Intervention	Study endpoints	Effects on study endpoints
Brzezinski et al. (2005)	Different populations including insomnia	17/284	Melatonin 0.3–40 mg versus placebo	SOL, TST, SE	SOL ↓; TST ↑; SE ↑
Buscemi et al. (2005)	Primary sleep disorders	14/425	Melatonin 1–5 mg versus placebo	SOL, WASO, TST, SE, SQ, USE	SOL ↓; best effect in sleep phase delay
Buscemi et al. (2006)	Secondary sleep disorders	15/524	Melatonin 1–10 mg versus placebo	SOL, USE	No effect on SOL, no USE
Braam et al. (2009)	Sleep problems with intellectual dysfunction	9/183	Melatonin 0.5–9 mg versus placebo	SOL, TST, NOA	SOL ↓; TST ↑; NOA ↑
Ferracioli-Oda et al. (2013)	Primary sleep disorders	19/1683	Melatonin 1—ca. 10 mg versus placebo	SOL, TST, SQ	Moderate effects on sleep continuity
Liu and Wang (2012)	Chronic insomnia	8/4055	Ramelteon 4–32 mg versus placebo	SOL, USE	Positive effects on subjective/objective SOL/no USE
McCleery et al. (2014)	Insomnia with Alzheimer's disease	5/313	Trazodon, melatonin, ramelteon	SOL, TST, WASO, SE	No evidence supporting melatonin/ramelteon
Kuriyama et al. (2014)	Insomnia	13/5812	Ramelteon	SOL, TST, SQ	SOL ↓; SQ ↑; clinically small effects
Winkler et al., 2014	Insomnia	31/3820	BZ, BZRA, sedating antidepressants, melatonin	PSG and subjective sleep parameters	Melatonin less effective than BZ/BZRA
Zhang et al. (2016)	Sleep disorders with neurodegenerative disorders	9/370	Melatonin	PSQI	Positive effects on PSQI and RBD
Auld et al. (2017)	Primary insomnia	5/1145	Melatonin	SOL	Significant effects for SOL
Marupuru et al. (2022)	Chronic insomnia > 60 years	17/2462	Melatonin, ramelteon	TST, SE, SOL, SQ	Significant effects for TST (PSG), SOL, SQ (subjective and PSG); no effect for SE

Abbreviations: BZ, benzodiazepines; BZRA, benzodiazepine receptor agonists; NOA, number of awakenings; PSG, polysomnography; PSQI, Pittsburgh Sleep Quality Index; RBD, rapid eye movement sleep behaviour disorder; SE, sleep efficiency; SOL, sleep-onset latency; SQ, sleep quality; TST, total sleep time; USE, undesired side-effects; WASO, wake time after sleep onset.

patients and patients with insomnia, in short-term studies. Several very recent all-encompassing (network) meta-analyses included studies on fast-release, PR melatonin and ramelteon (De Crescenzo et al., 2022; Hasan et al., 2023; Maruani et al., 2023; Yue et al., 2023); however, the results are conflicting. Considering this evidence, fast-release melatonin and ramelteon are not recommended for insomnia treatment (exception: if there are circadian factors involved) due to lack of efficacy. Given the evidence from studies with PR melatonin, and the fact that many health authorities in Europe have sanctioned its use beyond 4 weeks, PR melatonin, short- and longer-term administration can be considered (≥ 55 years.) after weighing the advantages and disadvantages.

A major caveat with melatonin is that it is manufactured by very different sources and can be purchased through a variety of avenues like supermarkets, internet, chemists and pharmacies. This probably results in marked variations in pharmacological quality. We thus urge caution and suggest users should obtain melatonin only through licensed pharmacies or through a prescription by their physician.

4.2.7 | Herbal remedies/phytotherapeutics

Table 11 lists herbal/phytotherapeutic substances available in Europe, and Table 15 presents all published meta-analyses on this topic, up until now.

Meta-analyses suggest small superiority of valerian compared with placebo (Leach & Page, 2015; Shinjyo et al., 2020). That said, this is in the context of rating the original studies to be of low quality. There is also evidence, also from low-quality original studies (Ji et al., 2021), that Saffron (Lian et al., 2022; Munirah et al., 2022), Viola odorata (Huang, Huang, et al., 2022) and diverse plant cocktails, from traditional Chinese medicine, may be effective to a low degree (Birling et al., 2020; Chen et al., 2022; Chen, Yi, et al., 2020; Fan et al., 2020; Li, Xu, et al., 2020; Lin et al., 2021; Wang et al., 2021; Wang, Ju, et al., 2020; Yang et al., 2019; Zhang, Liu, et al., 2019; Zhou et al., 2022). It is problematic for the authors of this guideline to evaluate these original studies, within the context of traditional Chinese medicine, as most of the original studies were not published in English language. Caution is therefore suggested given the conclusion by

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 15 Meta-analyses of phytotherapeutic interventions and their efficacy on the treatment of insomnia

Author (Year)	Population	No. studies/patients	Intervention	Study endpoints	Effects on study endpoints
Bent et al. (2006)	Insomnia	16/1093	Valerian versus placebo, short-term treatment	SQ, SOL	(a) Slight improvement in sleep quality (b) No improvement on other sleep parameters c) Poor quality of studies
Fernández-San-Martín et al. (2010)	Insomnia	18/1317	Valerian versus placebo	SQ	No effects on quantitative parameters, slight effects for SQ
Leach and Page (2015)	Insomnia	14/1602	Valerian, chamomile, kava, wuling	SOL, SE, TST, SQ	No significant effects
Ni et al. (2015)	Insomnia	76/7240	CHM versus placebo versus BZ	PSQI, CGI	CHM better than placebo, but poor quality studies
Khadivzadeh et al. (2018)	Insomnia (peri- and post-menopausal)	12/1257	Phytotherapeutics	Subjective parameters	Evidence considered as poor
Hieu et al. (2019)	Insomnia	1/34	Chamomilla	ISI	No evidence
Zhang, Liu, et al. (2019)	Primary insomnia	15/1500	CHM	PSQI, AIS, TST, SE, SOL	Significant effects for PSQI, AIS, TST, SE, SOL
Birling et al. (2020)	Insomnia disorder	19/1780	Zaoren Anshen (CHM)	PSQI, USE	Significant effects for Zaoren Anshen versus placebo; poor methodological quality studies
Chen, Yi, et al. (2020)	Insomnia	13/1175	Zaoren Anshen (CHM)	PSQI, USE	No evidence
Fan et al. (2020)	Insomnia disorder	13/1181	Long Dan Xie Gan Tang (CHM) versus BZ/BZRA/ barbiturate	Remission rate	Significant better remission with Long Dan Xie Gan Tang compared with BZ/BZRA/ barbiturates
Li, Xu, et al. (2020)	Insomnia disorder	14/1549	Yangxin Anshen (CHM) versus placebo/BZ/BZRA	PSQI, PSG parameter	Significant effects of Yangxin Anshen versus placebo for PSQI, PSG-parameter; no difference to BZ/BZRA
Shinjyo et al. (2020)	Diverse populations	60/6894	Valerian	Diverse outcomes	No clear effects
Wang et al. (2020)	Insomnia disorder	22/2029	Chaihu Longgu Muli (CHM)	PSQI, clinical effectiveness, USE	Significant effects of Chaihu Longgu on PSQI und clinical effectiveness; small USE
Hu et al. (2021)	Insomnia disorder + anxiety	9/681	Xiao Yao San (CHM)	PSQI	Significant effects of Xiao Yao San for PSQI
Ji et al. (2021)	Insomnia	72/7392	CHM	Various sleep parameters	Significant effects, very heterogenous quality of studies
Lin et al. (2021)	Insomnia disorder	14/910	Banxia (CHM)	PSQI	Significant effects for Banxia
Yoon et al. (2021)	Insomnia + cancer	14/1020	Phytotherapeutics	PSQI	Significant effects, low-quality studies
Huang, Huang, et al. (2022)	Insomnia	4/224	Viola odorata	SE, SOL, SQ, ISI, PSQI	Significant effects for SQ, ISI, PSQI; no effects for SE, SOL
Kim and Lim (2022)	Insomnia post-stroke	24/1942	Far eastern phytotherapeutics	PSQI	Significant effects on PSQI, low-quality studies
Lian et al. (2022)	Insomnia	8/611	Saffron	ISI, PSQI	Significant effects on ISI, PSQI

(Continues)

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 15 (Continued)

Author (Year)	Population	No. studies/patients	Intervention	Study endpoints	Effects on study endpoints
Luan et al. (2022)	Insomnia disorder	16/1260	Traditional chinese medicine	TST, SE, SOL, NOA, ISI, PSQI	Significant effects for many parameters, low-quality studies
Munirah et al. (2022)	Insomnia disorder	8/431	Crocus Sativus	TST, SQ, ISI	Significant effects for TST, SQ, ISI; low-quality studies
Zhou et al. (2022)	Insomnia disorder	19/1850	Wuling	PSQI	Significant effects on PSQI

Abbreviations: BZ, benzodiazepines; BZRA, benzodiazepine receptor agonists; CGI, clinical global impression; CHM, Chinese herbal medicine; ISI, Insomnia Severity Index; NOA, number of awakenings; PSG, polysomnography; PSQI, Pittsburgh Sleep Quality Index; SE, sleep efficiency; SOL, sleep-onset latency; SQ, sleep quality; TST, total sleep time; USE, undesired side-effects.

most authors of the meta-analyses that they consider the methodological quality of included studies as rather weak. As such, herbal/phytotherapeutic interventions cannot be recommended for either short- or long-term use to treat insomnia.

4.2.8 | Pharmacological long-term treatment

Studies on the pharmacotherapy of insomnia have mainly investigated treatment in the short term. Most studies on the BZs/BZRAs and low-dose sedating antidepressants had treatment durations shorter than 4 weeks, and the phase-III studies on daridorexant were for 3 months (with an extension up to 12 months). From a clinical perspective, given insomnia disorder is typically persistent, what would be more relevant would be to know the efficacy and safety of pharmacotherapy over a longer-term treatment period. It must be borne in mind, however, that all hypnotics are symptomatic treatments and insomnia symptoms typically return after discontinuation of these substances. Concerning longer-term treatment, therefore, up to this point, the scientific evidence remains insufficient to give a general positive recommendation for any medication beyond 4 or 12 weeks (depending on the type of drug being used). We do, however, acknowledge that in some cases, after carefully weighing the advantages and disadvantages, longer treatment periods might be initiated. Whether daridorexant or PR melatonin might become an exception to the rule remains to be seen, as more long-term data (in addition to practical experience) are needed. Thus, administration of these drugs for at least 3 months is an option, given that precautions should be taken to avoid the potential for tolerance and dependence.

4.3 | Comparative meta-analyses

The Institute of Quality and Economy in Health Care System (IQWiG) in Germany has prepared two evidence reports for the comparative treatment for insomnia in Germany. Both reports encompass meta-analytic summaries of the relevant literature.

The first report investigated internet- or mobile-based CBT-I (IQWiG-report—Nr. 1223; IQWIQ, 2021b), and compared efficacy with conventional CBT-I in adults with insomnia. Five studies were included in the report. With respect to the ISI, significant superiority for conventional CBT-I over dCBT-I was observed. This effect was measurable at the end of therapy and at follow-ups. The quality of evidence was considered, moderate.

The second report with the title "Cognitive Behavioural Therapy for Insomnia" (IQWiG-report—Nr. 1224; IQWIG, 2021a) investigated the efficacy of CBT-I compared with BZ and BZRA in adults with insomnia. Even 50 years after the introduction of both treatments, only five studies could be identified, making this comparison difficult. In some outcome parameters CBT-I was superior and in others pharmacological treatment was superior. The quality of evidence was rated as low for all outcomes, hindering valid conclusions. That said, a meta-analysis by Zhang et al. (2022) comparing CBT-I with any kind of pharmacological treatment for insomnia concluded that CBT-I may be superior to pharmacological treatment.

Recently a meta-analysis was published in the *Lancet*, comparing all pharmacological treatments for insomnia with each other (De Crescenzo et al., 2022). In this study, eszopiclone had the highest efficacy for insomnia-related parameters; however, this was explicitly tempered against the side-effects associated with this medication. Other drugs like BZ, doxepin and daridorexant were missing data on the efficacy in short- and long-term use according to the authors (De Crescenzo et al. (2022). Further, melatonin and other OTC medications were not considered effective according to this analysis.

4.4 | Daily use of medication versus as needed

With regard to sleep medications, it is mostly assumed that they are taken daily over short (up to 4 weeks) or longer-term (beyond 4 weeks) periods. This is probably engrained by the way clinical studies are conducted, mostly administering medication/placebo daily over a fixed time. Few studies have investigated intermittent use versus continuous use (Perlis et al., 2008), concluding that intermittent dosing does not lead to increased rebound phenomena. However, it could

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

not be determined whether intermittent dosing is equal or superior to continuous dosing.

Given that chronic insomnia (beyond 3 months) is characterised by fluctuations in sleep quality, the question of intermittent versus continuous medication intake is of high interest. In everyday clinical practice there is evidence that sleep medication is often prescribed/used on an "if needed" basis as opposed to a daily basis. The question now is how to define "as needed"—does this entail usage when somebody directly experiences sleep-onset problems or nocturnal awakening, or should the drug be taken prophylactically, i.e. the night before, when next-day challenges are expected that require a proper sleep beforehand. Depending on its half-life, drugs should not be taken too late in the night to avoid "hangover" problems, precluding middle of the night use.

Given these highly important questions, much more clinical routine data are needed to determine how to proceed. As of now, our recommendations refer to daily use (as informed by RCTs), but we do acknowledge that in daily clinical practice things might be handled very differently—so some patients may be long-term users over years or months, but may use one or two pills a week. Learning theory has made a strong point against intermittent use of potentially abusive substances, stating that intermittent positive reinforcement guarantees the strongest ties between a given substance and drug-taking behaviour (see overview by Perlis et al., 2008).

This issue becomes even more pertinent when we consider acute insomnia, an issue largely neglected by the insomnia field (with a few exceptions: see Ellis et al., 2012; Ellis, 2019). Our guideline refers to chronic insomnia (duration beyond 3 months) and we do not make any recommendations for acute/short-term insomnia. Nevertheless, in the medical arena, acute insomnia is a clinical issue facing many doctors routinely every day, for example, in acute situations like emergency visits or emergency admissions to hospitals or patients having to undergo impending life-saving procedures/surgery. We would assume that under these circumstances, swiftly acting medication may be very useful, according to the clinical picture and all ramifications of a given "case". We do assume that working on specific guidelines concerning these extreme scenarios will deserve special consideration and should be addressed in the near future.

4.5 | Tapering off medications

One major open question is whether CBT-I should be initiated when patients are on concomitant medication and request to be tapered off because they experience unpleasant side-effects, or feel they need support to discontinue the medication. For BZ/BZRA, specific schemata have been suggested with good success (Gould et al., 2014). The critical issue is whether all sleep medications should be tapered off prior to initiation of CBT-I or if an overlap can be accepted. There is evidence that CBT-I works better after discontinuation of sleep medication (Morin et al., 1994); however, in clinical practice, this may just not be feasible. It is suggested that those in the insomnia field work more closely together with experts in

TABLE 16 Recommendations for the treatment of chronic insomnia in adults of all ages

Treatment considerations
• Insomnia disorder should be actively treated whenever it presents (A)
• In the presence of comorbidities, clinical judgement should decide whether insomnia or the comorbid condition are treated first or whether both are treated at the same time (A)
CBT-I
• CBT-I should be provided as the the first-line treatment for insomnia disorder in adults of any age, regardless of comorbidities (A)
• CBT-I may be delivered either in-person or digitally (A)
• Sleep restriction and stimulus control are the most active ingredients of CBT-I (B)
Pharmacological interventions
• A pharmacological intervention can be proposed if CBT-I is not effective (A)
• BZs and BZRAs can be used in the short-term treatment of insomnia (≤ 4 weeks) (A)
• Longer-term treatment (off-label use) with BZ or BZRA, either daily or preferably intermittently, may be initiated in some cases, and the advantages and disadvantages need to be discussed on an individual basis (B)
• Low doses of sedating antidepressants can be considered (off-label use) in the short-term treatment of insomnia; contraindications have to be carefully considered (B)
• Longer-term treatment of insomnia disorder (without comorbidities; off-label use) with low-dose sedating antidepressants may be initiated in some cases, and the advantages and disadvantages need to be discussed on an individual basis (B)
• Orexin rexceptor antagonists can be used for a period of up to 3 months in the treatment of insomnia (A)
• Longer-term treatment of insomnia disorder with orexin receptor antagonists may be initiated in some cases, and the advantages and disadvantages need to be discussed on an individual basis (A)
• Because of insufficient evidence and possible risks, antihistamines are not recommended for insomnia treatment (A)
• Because of insufficient evidence and in light of their side-effects, antipsychotics are not recommended for insomnia treatment (A)
• Melatonin (fast-release, OTC or as a prescription drug) in general is not effective in the treatment of insomnia (A), if no circadian factors are involved
• Longer-term treatment of insomnia disorder with PR melatonin (in patients > 55 years) up to 3 months may be effective in some cases (B)
Other therapies
• Herbal remedies/phytotherapeutics are not recommended for the treatment of insomnia because of insufficient evidence (A). Light therapy and exercise regimes may be useful as adjunct therapies to CBT-I (B)

Abbreviations: BZ, benzodiazepines; BZRA, benzodiazepine receptor agonists; CBT-I, cognitive-behavioural therapy for insomnia; OTC, over-the-counter; PR, prolonged-release.

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License



FIGURE 1 Clinical algorithm. [Color figure can be viewed at wileyonlinelibrary.com]

addiction medicine to realise feasible solutions in this important area. Another important consideration is how to taper off any sleep medication if a patient is motivated to do so and explicitly requires to be helped. There may be cases of comorbid insomnia where one would want to argue against drug discontinuation, if there is expectancy of an extreme recurrence of insomnia symptoms during withdrawal that would negatively interfere with a medical comorbidity—for example, in cases of acute psychosis or mania.

5 | FINAL REMARKS ABOUT THE MANAGEMENT OF INSOMNIA AND FUTURE RESEARCH

This guideline tries to bring together state of the science evidence, generated over the last 6 years, to improve the management of chronic insomnia. The diagnostic recommendations have been summarised in Table 4, and the therapeutic recommendations are given in Table 16 and Figure 1.

Guidelines are attempts to reconcile the arenas of evidence-based medicine and clinical reality, between which there is usually a huge gap. With this guideline, we have tried, as we should, to collect and analyse the available evidence at its highest level, i.e. published meta-analyses. On the other hand, and as documented throughout this guideline, we are aware that many clinicians consider this aspect of their discipline as artificial or not rooted in clinical reality. However, RCTs on their own and/or summarised in meta-analyses still constitute the most powerful instruments to tease out what works and what does not. Individual clinical knowledge and intuition is important; however, so-called eminence-based medicine was disregarded for a reason!

Although there are new data available on the diagnosis and treatment of insomnia, there are still several areas in need of more intensive research.

The chapter on comparative meta-analyses (Section 3) clearly shows that there are only a few studies that directly compare different treatment options for insomnia. Although guidelines now universally favour CBT-I, it would be useful to have studies directly comparing CBT-I with pharmacotherapy. Moreover, pharmacological treatments have rarely been compared in high-quality randomised clinical studies. It is very difficult, therefore, to establish whether new medications are inferior, equal or superior to established treatments.

The chapter on longer-term pharmacological treatment (Section 2.8) highlights a distinct need for longer-term treatment studies of insomnia using pharmacological agents. This lack of evidence has resulted in continued cautioning when considering pharmacotherapy in the long-term treatment of insomnia. We see a certain potential for PR melatonin and orexin-receptor-antagonists under these circumstances, especially in those patients for whom CBT-I was not effective.

An important question when judging treatment options for insomnia is also the question of placebo effects. Placebos have been shown to have small effects on subjective insomnia symptoms and even medium effects on global measures of sleep quality (Yeung et al., 2018). As such, the use of adequate placebos is a very important methodological

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

precondition to judge the efficacy of treatment options in insomnia. Presently, there is a paucity of appropriate placebo-controlled studies with CBT-I, with the majority of studies using waiting list or treatment as usual control conditions. That said, Van Straten et al. (2018) compared waitlist CBT-I studies with placebo-controlled CBT-I studies, and concluded that the effects were comparable. It also has been demonstrated that Sleepio (one of the available dCBT-I platforms) is superior to placebo (Espie et al., 2012). Future clinical insomnia research on non-pharmacological treatments should therefore pay greater attention to the fact that it is feasible to construct valid placebo conditions. Additionally, the effects of CBT-I on objectively measured sleep parameters, from PSG, remain under-researched (Mitchell et al., 2019).

Another important issue concerns the question of remission versus response/non-response in both CBT-I and pharmacological trials. Morin et al. (2020) showed that in a comparative trial of psychological therapies versus zolpidem, clinically meaningful response occurred in 45% of psychological therapies (remission in slightly less than 40%) and in 49% of zolpidem-treated subjects (remission in slightly less than 30%). Most studies in the field (concerning both CBT-I and pharmacotherapy) usually report pre- to post-comparisons of outcome variables and effect strengths. Future research will have to take a closer look at response versus remission rates and present strategies to achieve higher remission rates, as in studies with sequential designs such as the one carried out by Morin et al. (2020).

The increasing trials pipeline on single-component CBT-I interventions is to be encouraged (Kyle et al., 2023), both with respect to the assessment of mechanisms and their unique efficacy. Recognition that CBT-I is an amalgam of numerous cognitive-behavioural options, each of which may have a discrete evidence base, offers considerable promise, and further research on potentially differing dimensions or sequencing of behavioural and cognitive therapeutics and their comparative effectiveness appears to be warranted.

A strong emphasis should be put on researching combination therapies, especially between medications and psychotherapeutic approaches. We need to know whether patients should be tapered off medication prior to CBT-I or not—will CBT-I then work better? Might a combination of medication and CBT-I work better in some patients or not?

What about patients who are dependent on hypnotic medication—how can we help them if they are motivated to abstain? Tapering strategies need to be developed and integrated with CBT-I strategies.

Further treatment options, which currently lack a solid empirical basis, need to be taken account of in future investigations, such as the cannabinoids (Bhagavan et al., 2020), and tES or TMS (Krone et al., 2023).

The treatment of insomnia and comorbid medical disorders is still under-researched. Attention should be drawn for example to "comorbid insomnia and sleep apnea" (COMISA). Some initial evidence seems to support CBT-I alongside treatment of sleep apnea, but further trials would be welcome.

A further research step will have to include the practical implementation of any kind of insomnia management in everyday clinical practice. Our knowledge is primarily based on good-quality evidence from RCTs; however, it is well known that high-quality RCTs may be very far from what happens in clinical routine. Thus, we need data from everyday clinical practice about diagnostic procedures that are accepted and executed, treatment acceptance, adherence, attrition, side-effects and therapeutic efficacy. For example, pragmatic adaptations of CBT-I for patients with insomnia in acute psychiatric care are currently developed and very promising (Schneider et al., 2020).

6 | GUIDELINE RECOMMENDATIONS AND CLINICAL ALGORITHM

Recommendations for the diagnostic management of chronic insomnia and its comorbidities in adults of all ages have been summarised in Table 4. Recommendations for insomnia treatment are shown in Table 16.

A clinical algorithm summarising the diagnostic and therapeutic process is shown in Figure 1.

AUTHOR CONTRIBUTIONS

Dieter Riemann: Conceptualization; methodology; data curation; supervision; project administration; writing – original draft; writing – review and editing; validation; formal analysis; visualization; resources. **Colin A. Espie:** Conceptualization; methodology; writing – original draft; writing – review and editing; project administration. **Ellemarije Altena:** Conceptualization; writing – original draft; writing – review and editing. **Erna Sif Arnardottir:** Conceptualization; writing – original draft; writing – review and editing; project administration. **Chiara Baglioni:** Conceptualization; methodology; writing – original draft; writing – review and editing. **Claudio L. A. Bassetti:** Conceptualization; writing – original draft; writing – review and editing. **Celyne Bastien:** Conceptualization; writing – original draft; writing – review and editing. **Natalija Berzina:** Conceptualization; writing – original draft; writing – review and editing. **Bjørn Bjorvatn:** Conceptualization; methodology; writing – original draft; writing – review and editing. **Dimitris Dikeos:** Conceptualization; writing – original draft; writing – review and editing. **Leja Dolenc Groselj:** Conceptualization; writing – original draft. **Jason G. Ellis:** Conceptualization; methodology; writing – original draft; writing – review and editing. **Diego Garcia-Borreguero:** Conceptualization; methodology; writing – original draft; writing – review and editing. **Pierre Geoffroy A:** Conceptualization; writing – original draft; writing – review and editing. **Michaela Gjerstad:** Conceptualization; writing – original draft; writing – review and editing. **Marta Gonçalves:** Conceptualization; writing – original draft; writing – review and editing. **Elisabeth Hertenstein:** Conceptualization; methodology; writing – original draft; writing – review and editing. **Kerstin Hoedlmoser:** Conceptualization; writing – original draft; writing – review and editing. **Tuuliki Hion:** Conceptualization; writing – original draft; writing – review and editing. **Brigitte Holzinger:** Conceptualization; writing – original draft; writing – review and editing. **Karolina Janku:** Conceptualization; writing – original draft; writing – review and editing. **Markus Jansson-Fröjmark:** Conceptualization; writing – original draft; writing – review and editing. **Heli Järnefelt:** Conceptualization; writing – original draft; writing – review and editing. **Susanna Jernelöv:** Conceptualization; methodology; writing – original draft; writing – review

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

and editing. **Poul Jørgen Jennum:** Conceptualization; writing – original draft; writing – review and editing. **Samson Khachatryan:** Conceptualization; writing – original draft; writing – review and editing. **Lukas Krone:** Conceptualization; methodology; writing – original draft; writing – review and editing. **Simon D. Kyle:** Conceptualization; methodology; writing – original draft; writing – review and editing. **Jaap Lancee:** Conceptualization; writing – original draft; writing – review and editing. **Damien Leger:** Conceptualization; writing – original draft; writing – review and editing. **Adrian Lupusor:** Conceptualization; writing – original draft; writing – review and editing. **Daniel Ruivo Marques:** Conceptualization; writing – original draft; writing – review and editing. **Christoph Nissen:** Conceptualization; methodology; supervision; writing – original draft; writing – review and editing. **Laura Palagini:** Conceptualization; supervision; writing – original draft; writing – review and editing. **Tiina Paunio:** Conceptualization; writing – original draft; writing – review and editing. **Lampros Perogamvros:** Conceptualization; writing – original draft; writing – review and editing. **Dirk Pevernagie:** Conceptualization; methodology; writing – original draft; writing – review and editing. **Manuel Schabus:** Conceptualization; writing – original draft; writing – review and editing. **Tamar Shochat:** Conceptualization; writing – original draft; writing – review and editing. **Andras Szentkiralyi:** Conceptualization; writing – original draft; writing – review and editing. **Eus Van Someren:** Conceptualization; methodology; supervision; writing – original draft; writing – review and editing. **Annemieke van Straten:** Conceptualization; writing – original draft; writing – review and editing. **Adam Wichniak:** Conceptualization; writing – original draft; writing – review and editing. **Kai Spiegelhalder:** Conceptualization; methodology; data curation; supervision; formal analysis; writing – original draft; writing – review and editing; project administration.

AFFILIATIONS

[1]Department of Clinical Psychology and Psychophysiology, Centre for Mental Health (Department), Medical Center – University of Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany
[2]Center for Basics in NeuroModulation (NeuroModulBasics), Faculty of Medicine, University of Freiburg, Freiburg, Germany
[3]Sleep and Circadian Neuroscience Institute, Nuffield Department of Clinical Neuroscience at the University of Oxford, Oxford, UK
[4]Université de Bordeaux, Bordeaux, France
[5]Reykjavik University Sleep Institute, School of Technology, Reykjavik University, Reykjavik, Iceland
[6]Landspitali – The National University Hospital of Iceland, Reykjavik, Iceland
[7]Human Sciences Department, University of Rome Guglielmo Marconi Rome, Rome, Italy
[8]Department of Neurology, Inselspital, University of Bern, Berne, Switzerland
[9]École de Psychologie, Université Laval, Québec, Québec, Canada
[10]Psychiatric Practice, Riga, Latvia
[11]Department of Global Public Health and Primary Care, University of Bergen, Bergen, Norway
[12]First Department of Psychiatry, Medical School, National and Kapodistrian University of Athens, Athens, Greece
[13]Institute of Clinical Neurophysiology, University Medical Center Ljubljana, Ljubljana, Slovenia
[14]Northumbria Sleep Research Laboratory, Northumbria University, Newcastle, UK
[15]Sleep Research Institute, Madrid, Spain
[16]Department of Psychiatry, Paris Cite University, Paris, France
[17]Stavanger University Hospital, Stavanger, Norway
[18]Centro de Medicina de Sono, Hospital Cuf, Porto, Portugal
[19]University Hospital of Psychiatry and Psychotherapy, University of Bern, Bern, Switzerland
[20]Centre for Cognitive Neurosciences, University of Salzburg, Salzburg, Austria
[21]East-Viru Central Hospital, Kohtla-Järve, Estonia
[22]Institute for Consciousness and Dream Research, Vienna, Austria
[23]Center for Sleep and Chronobiology Research, National Institute of Mental Health, Klecany, Czech Republic
[24]Division of Psychology, Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden
[25]Centre for Psychiatry Research, Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden and Stockholm Health Care Services, Stockholm, Sweden
[26]Finnish Institute of Occupational Health, Helsinki, Finland
[27]Department of Clinical Medicine, University of Copenhagen, Kobenhavn, Denmark
[28]Department of Neurology and Neurosurgery, Armenian National Institute of Health, Yerevan, Armenia
[29]Department of Clinical Psychology, University of Amsterdam, Amsterdam, The Netherlands
[30]Université Paris Cité, APHP, Hôtel Dieu de Paris, Centre du Sommeil et de la Vigilance, Paris, France
[31]Functional Neurology, Institute of Neurology and Neurosurgery, Chisinau, Moldova
[32]Department of Education and Psychology, University of Aveiro, Aveiro, Portugal
[33]CINEICC – Center for Research in Neuropsychology and Cognitive Behavioral Intervention, Faculty of Psychology and Educational Sciences, University of Coimbra, Coimbra, Portugal
[34]Department of Psychiatry, University Hospital Geneve, Geneve, Switzerland
[35]Psychiatry Unit, Department of Clinical and Experimental Medicine, School of Medicine, University of Pisa, Pisa, Italy
[36]Department of Psychiatry and SleepWell Research Program, Faculty of Medicine, University of Helsinki and Helsinki University Central Hospital, Helsinki, Finland; Public Health and Welfare, Finnish Institute for Health and Welfare, Helsinki, Finland
[37]Department of Internal Medicine and Pediatrics, Ghent University, Ghent, Belgium
[38]The Cheryl Spencer Institute of Nursing Research, University of Haifa, Haifa, Israel
[39]Institute of Epidemiology and Social Medicine, University of Münster, Münster, Germany

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

[40]Department of Sleep and Cognition, Netherlands Institute for Neuroscience, Royal Netherlands Academy of Arts and Sciences, Amsterdam, The Netherlands

[41]Departments of Integrative Neurophysiology and Psychiatry, Center for Neurogenomics and Cognitive Research, Amsterdam UMC, Amsterdam Neuroscience, VU University, Amsterdam, The Netherlands

[42]Department of Clinical, Neuro- and Developmental Psychology & Amsterdam Public Health Research Institute, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

[43]Sleep Medicine Center and Third Department of Psychiatry, Institute of Psychiatry and Neurology, Warsaw, Poland

[44]Multidisciplinary Sleep Disorders Centre, Antwerp University Hospital and University of Antwerp, Antwerp, Belgium

ACKNOWLEDGEMENTS

This guideline was supported and endorsed by the European Sleep Research Society (ESRS) and by the European Insomnia Network. We would like to thank especially the board of the ESRS in its present composition for having the confidence to trust us in delegating us to write this guideline. The ESRS generously provided € 1000 for financing English proof reading prior to JSR submission. Beyond, no financial support was available. Open Access funding enabled and organized by Projekt DEAL.

CONFLICT OF INTEREST STATEMENT

All conflict of interests forms are available as supplemental material.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Dieter Riemann https://orcid.org/0000-0002-1968-6220
Colin A. Espie https://orcid.org/0000-0002-1294-8734
Erna Sif Arnardottir https://orcid.org/0000-0003-0877-3529
Chiara Baglioni https://orcid.org/0000-0003-3150-7755
Leja Dolenc Groselj https://orcid.org/0000-0002-8350-951X
Jason G. Ellis https://orcid.org/0000-0002-8496-520X
Pierre A. Geoffroy https://orcid.org/0000-0001-9121-209X
Karolina Janku https://orcid.org/0000-0002-6804-6967
Markus Jansson-Fröjmark https://orcid.org/0000-0003-2059-1621
Susanna Jernelöv https://orcid.org/0000-0002-0633-8104
Samson Khachatryan https://orcid.org/0000-0002-3098-2135
Lukas Krone https://orcid.org/0000-0002-5535-7221
Jaap Lancee https://orcid.org/0000-0002-7722-5780
Damien Leger https://orcid.org/0000-0003-1168-480X
Daniel Ruivo Marques https://orcid.org/0000-0003-3729-0120
Christoph Nissen https://orcid.org/0000-0001-9809-0275
Laura Palagini https://orcid.org/0000-0003-1676-629X
Dirk Pevernagie https://orcid.org/0000-0002-7372-8583
Tamar Shochat https://orcid.org/0000-0002-4371-3567
Eus Van Someren https://orcid.org/0000-0002-9970-8791
Annemieke van Straten https://orcid.org/0000-0001-6875-2215
Kai Spiegelhalder https://orcid.org/0000-0002-4133-3464

REFERENCES

AASM (American Academy of Sleep Medicine). (2014). International classification of sleep disorders, 3rd edition. In *American Academy of sleep medicine (editor)*. Rochester, MN.

Abetz, L., Arbuckle, R., Allen, R. P., Garcia-Borreguero, D., Hening, W., Walters, A., Mavraki, E., & Kirsch, J. M. (2006). The reliability, validity and responsiveness of the IRLSSG rating scale and subscales in a clinical-trial setting. *Sleep Medicine*, *4*, 340–349.

Alimoradi, Z., Jafari, E., Broström, A., Ohayon, M. M., Lin, C. Y., Griffiths, M. D., Blom, K., Jernelöv, S., Kaldo, V., & Pakpour, A. H. (2022). Effects of cognitive behavioral therapy for insomnia (CBT-I) on quality of life: A systematic review and meta-analysis. *Sleep Medicine Reviews*, *64*, 101646. https://doi.org/10.1016/j.smrv.2022.101646

Amiri, S., Hasani, J., & Satkin, M. (2021). Effect of exercise training on improving sleep disturbances: A systematic review and meta-analysis of randomized control trials. *Sleep Medicine*, *84*, 205–218.

APA (American Psychiatric Association). (2013). Diagnostic and statistical manual of mental disorders (5th edition: DSM-5). In *American Psychiatric Association (editor)*. DC.

Auld, F., Maschauer, E. L., Morrison, I., Skene, D. J., & Riha, R. L. (2017). Evidence for the efficacy of melatonin in the treatment of primary adult sleep disorders. *Sleep Medicine Reviews*, *34*, 10–22.

Baglioni, C., Espie, C. A., & Riemann, D. (2022). *Cognitive-behavioural therapy for insomnia (CBT-I) across the lifespan*. Guidelines and clinical protocols for health professionals. Wiley & Sons.

Baglioni, C., Altena, E., Bjorvatn, B., Blom, K., Bothelius, K., Devoto, A., Espie, C. A., Frase, L., Gavriloff, D., Hion, T., Hoflehner, A., Högl, B., Holzinger, B., Järnefelt, H., Jernelöv, S., Johann, A. F., Lombardo, C., Nissen, C., Palagini, L., … Riemann, D. (2020). The european academy for cognitive behavioural therapy for insomnia: An initiative of the european insomnia network to promote implementation and dissemination of treatment. *Journal of Sleep Research*, *29*, e12967. https://doi.org/10.1111/jsr.12967

Baglioni, C., Battagliese, G., Feige, B., Spiegelhalder, K., Nissen, C., Voderholzer, U., Lombardo, C., & Riemann, D. (2011). Insomnia is a predictor of depression: A meta-analytic evaluation of longitudinal epidemiological studies. *Journal of Affective Disorders*, *135*, 10–19.

Baglioni, C., Bostanova, Z., Bacaro, V., Benz, F., Hertenstein, E., Spiegelhalder, K., Rücker, G., Frase, L., Riemann, D., & Feige, B. (2020). A systematic review and network meta-analysis of randomized controlled trials evaluating the evidence base of melatonin, light exposure, exercise, and complementary and alternative medicine for patients with insomnia disorder. *Journal of Clinical Medicine*, *9*(6), 1949. https://doi.org/10.3390/jcm9061949

Baglioni, C., Regen, W., Teghen, A., Spiegelhalder, K., Feige, B., Nissen, C., & Riemann, D. (2014). Sleep changes in the disorder of insomnia: A meta-analysis of polysomnographic studies. *Sleep Medicine Reviews*, *18*, 195–213.

Baka, A., van der Zweerde, T., Lancee, J., Bosmans, J. E., & van Straten, A. (2022). Cost-effectiveness of guided internet delivered cognitive behavioral tcerapy in comparison with care-as-usual for patients with insomnia in general practice. *Behavioral Sleep Medicine*, *20*(2), 188–203.

Ballesio, A., Aquino, M., Feige, B., Johann, A. F., Kyle, S. D., Spiegelhalder, K., Lombardo, C., Rücker, G., Riemann, D., & Baglioni, C. (2018). The effectiveness and behavioural and cognitive therapies for insomnia on depressive and fatigue symptoms: A systematic review and network meta-analysis. *Sleep Medicine Reviews*, *37*, 114–129.

Ballesio, A., Bacaro, V., Vacca, M., Chirico, A., Lucidi, F., Riemann, D., Baglioni, C., & Lombardo, C. (2021). Does cognitive behaviour therapy

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

for insomnia reduce repetitive negative thinking and sleep-related worry beliefs? A systematic review and meta-analysis. *Sleep Medicine Reviews*, *55*, 101378. https://doi.org/10.1016/j.smrv.2020.101378

Banno, M., Harada, Y., Taniguchi, M., Tobita, R., Tsujimoto, H., Tsuhimoto, Y., Kataoka, Y., & Noda, A. (2018). Exercise can improve sleep quality: A systematic review and meta-analysis. *PeerJournal*, *6*, e5172.

Barker, M. J., Greenwood, K. M., Jackson, M., & Crowe, S. F. (2004). Persistence of cognitive effects after withdrawal from long-term benzodiazepine use: A meta-analysis. *Archives of Clinical Neuropsychology*, *19*, 437–454.

Bassetti, C., Ferini-Strambe, L., Adamantidis, A., Benedetti, F., Brown, S., Bruni, C. O., Cajochen, C., Dolenc-Groselj, L., Ferri, R., Gais, R., Huber, R., Khatami, R., Lammers, G., Manconi, M., Nissen, C., Nobili, L., Peigneux, P., Pollmächer, T., Randerath, W., … Wetter, T. P. (2015). Neurology and psychiatry: Waking up to the opportunities of sleep. *European Journal of Neurology*, *10*, 1337–1354.

Bassetti, C., Randerath, W., Vignatelli, L., Ferini-Strambe, L., Brill, A. K., Bonsignore, M. R., Groze, L., Jennum, P., Minnerup, J., Nobili, L., Tonia, T., Morgan, R., Kerry, J., Riha, R., McNicholas, W., & Papavasileiou, V. (2020). EAN/ESO/ESRS statement on the impact of sleep and sleep disorders on risk and outcome of stroke. *European Journal of Neurology*, *27*(7), 1117–1136. https://doi.org/10.1111/ene.14201

Bastien, C. H., Vallières, A., & Morin, C. M. (2001). Validation of the insomnia severity index as an outcome measure for insomnia research. *Sleep Medicine*, *2*, 297–307.

Behrendt, D., Ebert, D. D., Spiegelhalder, K., & Lehr, D. (2020). Recovery training in improving sleep in workers: Randomized controlled trial in the general working population. *Journal of Medical Internet Research*, *22*(1), e13346. https://doi.org/10.2196/13346

Belleville, G., Cousineau, H., Levrier, K., & St. Pierre-Delorme, M. E. (2011). Meta-analytic review of the impact of cognitive-behavior therapy for insomnia on concomitant anxiety. *Clinical Psychology Reviews*, *31*, 638–652.

Bent, S., Padula, A., Moore, D., Patterson, M., & Mehling, W. (2006). Valerian for sleep: A systematic review and meta-analysis. *American Journal of Medicine*, *119*, 1005–1012.

Benz, F., Knoop, T., Ballesio, A., Bacaro, V., Johann, A. F., Rücker, G., Feige, B., Riemann, D., & Baglioni, C. (2020). The efficacy of cognitive and behavior therapies for insomnia on daytime symptoms: A systematic review and network meta-analysis. *Clinical Psychology Reviews*, *80*, 101873.

Bhagavan, C., Kung, S., Doppen, M., John, M., Vakalalabure, I., Oldfield, K., Braithwaite, I., & Newton-Howes, G. (2020). Cannabinoids in the treatment of insomnia disorder: A systematic review and meta-analysis. *CNS Drugs*, *34*, 1217–1228.

Birling, Y., Jia, M., Li, G., Sarris, J., Bensoussan, A., & Zhu, X. (2020). Zao Ren an Shen for insomnia: A systematic review with meta-analysis. *Sleep Medicine*, *69*, 41–50.

Bisdounis, L., Saunders, K. E. A., Farley, H. J., Lee, C. K., McGowan, N. M., Espie, C. A., & Kyle, S. D. (2022). Psychological and behavioural interventions in bipolar disorder that target sleep and circadian rhythms: A systematic review of randomized controlled trials. *Neuroscience and Biobehavioral Reviews*, *132*, 378–390.

Bjorvatn, B., Fiske, E., & Pallesen, S. (2011). A self-help book is better than sleep hygiene advice for insomnia: A randomized controlled comparative study. *Scandinavian Journal of Psychology*, *52*, 580–585.

Blom, K., Jernelov, S., Ruck, C., Lindefors, N., & Kaldo, V. (2016). Three-year follow-up of insomnia and hypnotics after controlled internet treatment for insomnia. *Sleep*, *39*(6), 1267–1274.

Bomhof-Roordink, H., Gärtner, F. R., Stiggelbout, A. M., & Pieterse, A. H. (2019). Key components of shared decison making models: A systematic review. *British Medical Journal Open*, *9*, e031763.

Bootzin, R. (1972). A stimulus control treatment for insomnia. *Proceedings of the American Psychological Association*, *7*, 395–396.

Borbély, A. A. (1982). A two process model of sleep regulation. *Human Neurobiology*, *1*, 195–204.

Braam, W., Smits, M. G., Didden, R., Korzilius, H., van Geijlswijk, I. M., & Curfs, L. M. G. (2009). Exogenous melatonin for sleep problems in individuals with intellectual disability: A meta-analysis. *Developmental Medicine & Child Neurology*, *51*, 340–348.

Brzezinski, A., Vangel, M. G., Wurtman, R. J., Norrie, G., Zhdanova, I., Ben-Shushan, A., & Ford, I. (2005). Effects of exogenous melatonin on sleep: A meta-analysis. *Sleep Medicine Reviews*, *9*, 41–50.

Buscemi, N., Vandermeer, B., Friesen, C., Bialy, L., Tubman, M., Ospina, M., Klassen, T. P., & Witmans, M. (2007). The efficacy and safety of drug treatments for chronic insomnia in adults: A meta-analysis of RCTs. *Journal of General Internal Medicine*, *22*, 1335–1350.

Buscemi, N., Vandermeer, B., Hootin, N., Pandya, R., Tjosvold, L., Hartling, L., Vohra, S., Klassen, T. P., & Baker, G. (2006). Efficacy and safety of exogenous melatonin for secondary sleep disorders and sleep disorders accompanying sleep restriction: Meta-analysis. *British Medical Journal*, *332*, 385–393.

Buscemi, N., Vandermeer, B., Hooton, N., Pandya, R., Tjosvold, L., Hartling, L., Baker, G., Klassen, T. P., & Vohra, S. (2005). The efficacy and safety of exogenous melatonin for primary sleep disorders. A meta-analysis. *Journal of General Internal Medicine*, *20*, 1151–1158.

Buysse, D. J., Reynolds, C. F., 3rd, Monk, T. H., Berman, S. R., & Kupfer, D. J. (1989). The Pittsburgh sleep quality index: A new instrument for psychiatric practice and research. *Psychiatry Research*, *28*, 193–213.

Carney, C. E., Buysse, D. J., Ancoli-Israel, S., Edinger, J. D., Krystal, A. D., Lichstein, K., & Morin, C. M. (2012). The consensus sleep diary: Standardizing prospective sleep self-monitoring. *Sleep*, *35*, 287–302.

Chambe, J., Reynaud, E., Maruani, J., Fraih, E., Geoffroy, P. A., & Bourgin, P. (2023). Light therapy in insomnia disorder: A systematic review and meta-analysis. *Journal of Sleep Research*, *2023*, e13895. https://doi.org/10.1111/jsr.13895

Chapoutot, M., Peter-Derex, L., Schoendorff, B., Faivre, T., Bastuji, H., & Putois, B. (2021). Telehealth-delivered CBT-I programme enhanced by acceptance and commitment therapy for insomnia and hypnotic dependence: A pilot randomized controlled trial. *Journal of Sleep Research*, *30*(1), e13199. https://doi.org/10.1111/jsr.13199

Chen, B. W., Yi, J., Sun, B., Jia, P., Zhang, W. J., & Liu, B. Y. (2020). Efficacy and safety of Zaoren Anshen capsules in the treatment of insomnia: A meta-analysis of randomized controlled trials. *Medicine (Baltimore)*, *99*(6), e19033.

Chen, C. T., Tung, H. H., Fang, C. J., Wang, J. L., Ko, N. Y., Chang, Y. J., & Chen, Y. C. (2021). Effect of music therapy on improving sleep quality in older adults: A systematic review and meta-analysis. *Journal of the American Geriatric Society*, *69*, 1925–1932.

Chen, F., Chen, Z., Cheng, Y., Li, J., Liao, R., Zhao, Z., Wu, K., & Liu, J. (2022). Meta analysis for insomnia Guizhi Gancao Longgu Muli decoction for insomnia a meta-analysis. *Complementary Therapy Clinical Practice*, *47*, 101550.

Chen, T. L., Chang, S. C., Hsieh, H. F., Huang, C. Y., Chuang, J. H., & Wang, H. H. (2020). Effects of mindfulness-based stress reduction on sleep quality and mental health for insomnia patients: A meta-analysis. *Journal of Psychosomatic Research*, *135*, 10144.

Cheng, S. K., & Dizon, J. (2012). Computerised cognitive behavioural therapy for insomnia: A systematic review and meta-analysis. *Psychotherapy and Psychosomatics*, *81*, 206–216.

Choong, S. Y. X., Tan, X. Y. J., Cheng, L. J., & Lau, Y. (2022). Effectiveness of exercise in improving sleep outcomes among perinatal women: A systematic review and meta-analysis of randomised controlled trials. *Behavioral Sleep Medicine*, *20*, 410–428.

Chung, F., Yegneswaran, B., Liao, P., Vairaanathan, S., Islam, S., Khajedehi, A., & Shapiro, C. (2008). STOP questionnaire – A tool to screen patients for obstructive sleep apnea. *Anesthesiology*, *108*, 812–821.

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Chung, K. F., Lee, C. T., Yeung, W. F., Chan, M. S., Chung, E. W., & Lin, W. L. (2018). Sleep hygiene education as a treatment of insomnia: A systematic review and meta-analysis. *Family Practice*, *35*, 365–375.

Condon, H. E., maurer, L. F., & kyle, S. D. (2021). Reporting adverse events in cognitive behavioral therapy for insomnia: A systematic examination of randosmised controlled trials. *Sleep Medicine Reviews*, *56*, 101412.

Crönlein, T., Geisler, P., Langguth, B., Eichhammer, P., Jara, C., Pieh, C., Zulley, J., & Hajak, G. (2012). Polysomnography reveals unexpectedly high rates of organic sleep disorders in patients with prediagnosed primary insomnia. *Sleep and Breathing*, *16*, 1097–1103.

Curtis, F., Laparidou, D., Bridle, C., Law, G. R., Durrant, S., Rodriguez, A., Pierzycki, R. H., & Siriwardena, A. N. (2021). Effects of cognitive behavioural therapy on insomnia in adults with tinnitus: Systematic review and meta-analysis of randomised controlled trials. *Sleep Medicine Reviews*, *56*, 1405.

Damsgard, L., Janbek, J., Laursen, T. M., Erlangsen, A., Spira, A. P., & Waldemar, G. (2022). Hospital-diagnsoed sleep disorders and incident dementia: A nationwide observational cohort study. *European Journal of Neurology*, *29*, 3528–3536.

D'Aurea, C. V. R., Frange, C., Poyares, D., Souza, A. A. L., & Lenza, M. (2022). Physical exercise as a therapeutic approach for adults with insomnia: Systematic review and meta-analysis. *Einstein (Sao Paolo)*, *20*, eAO8058.

De Crescenzo, F., D'Alò, G. L., Ostinelli, E. G., Ciabattini, M., Di Franco, V., Watanabe, N., Kurtulmus, A., Tomlinson, A., Mitrova, Z., Foti, F., Del Giovane, C., Quested, D. J., Cowen, P. J., Barbui, C., Amato, L., Efthimiou, O., & Cipriani, A. (2022). Comparative effects of pharmacological interventions for the acute and long-term management of insomnia disorder in adults: A systematic review and network meta-analysis. *Lancet*, *400*, 170–184.

Dekker, K., Benjamins, J. S., Maksimovic, T., Filardi, M., Hofman, W. F., van Straten, A., & Van Someren, E. J. W. (2020). Combined internet-based cognitive-behavioral and chronobiological intervention for insomnia: A randomized controlled trial. *Psychotherapy and Psychosomatics*, *89*, 117–118.

Dikeos, D., Wichniak, A., Ktonas, P. Y., Mikoteit, T., Crönlein, T., Eckert, A., Koprivova, J., Ntafouli, M., Spiegelhalder, K., Hatzinger, M., Riemann, D., & Soldatos, C. (2023). The potential of biomarkers for diagnosing insomnia: Consensus statement of the WFSBP task force on sleep disorders. *World Journal of Biological Psychiatry*, *7*, 1–29. https://doi.org/10.1080/15622975.2023.2171479

Dressle, R., & Riemann, D. (2023). Hyperarousal in insomnia disorder: Current evidence and potential mechanisms. *Journal of Sleep Research*, *32* (6), 13928. https://doi.org/10.1111/jsr.13928

Dündar, Y., Dodd, S., Strobl, J., Boland, A., Dickson, R., & Walley, T. (2004). Comparative efficacy of newer hypnotic drugs for the short-term management of insomnia: A systematic review and meta-analysis. *Human Psychopharmacology*, *19*, 305–322.

Dupuy, L., Morin, C. M., de Sevin, E., Taillard, J., Salles, N., Bioulac, S., Auriacombe, M., Micoulaud-Franchi, J. A., & Philip, P. (2022). Smartphone-based virtual agents and insomnia management: A proof-of-concept study for new methods of autonomous screening and management of insomnia symptoms in the general population. *Journal of Sleep Research*, *31*(2), e13489.

Ebert, D. D., Berking, M., Thiart, H., Riper, H., Laferton, J. A. C., Cuijpers, P., Sieland, B., & Lehr, D. (2015). Restoring depleted resources: Efficacy and mechanisms of change of an internet-based unguided recovery training for better sleep and psychological detachment from work. *Health Psychology*, *34S*, 1240–1251. https://doi.org/10.1037/hea0000277

Edinger, J. D., Arnedt, J. T., Bertisch, S. M., Carney, C. E., Harrington, J. J., Lichstein, K. L., Sateia, M. J., Troxel, W. M., Zhou, E. S., Kazmi, U., Heald, J. L., & Martin, J. L. (2021). Behavioral and psychological treatments for chronic insomnia disorder in adults: An American Academy of sleep medicine clinical practice guideline. *Journal of Clinical Sleep Medicine*, *17*, 255–262.

Eigl, E. S., Hauser, T., & Schabus, M. (2022). On the efficacy of a web-based treatment program for sleep disorders and insomnia – a pilot study. Poster presented at the 26th congress of the European Sleep Research Society (ESRS), Athens, Greece, September 27th – 30th 2022.

El Rafihi-Ferreira, R., Morin, C. M., Toscanini, A. C., Lotufo, N., Brasil, I. S., Gallinaro, J. G., Borges, D. S., Conway, S. G., & Hasan, R. (2021). Acceptance and commitment therapy-based behavioral intervention for insomnia: A pilot randomized controlled trial. *Brazilian Journal of Psychiatry*, *43*, 504–509.

Ellis, J., Gehrman, P., Espie, C., Riemann, D., & Perlis, M. (2012). Acute insomnia: current conceptualizations and future directions. *Sleep Medicine Reviews*, *16*, 5–14.

Ellis, J. G. (2019). Cognitive behavioral therapy for insomnia and acute insomnia: Considerations and controversies. *Sleep Medicine Clinics*, *14*(2), 267–274.

El-Rafihi-Ferreira, R., Morin, C. M., Hasan, R., Brasil, I. S., Zago-Ribeiro, J. H., & Cecilia Toscanini, A. (2022). A pilot randomized controlled trial (RCT) of acceptance and commitment therapy versus cognitive behavioral therapy for chronic insomnia. *Behavioral Sleep Medicine*, *10*, 1–15.

EMA (European Medicines Agency). (2007/2010/2023). Circadin (melatonin). https://www.ema.europa.eu/en/medicines/human/EPAR/Circadin

EMA (European Medicines Agency). (2022/2023). Quvivig (daridorexant). https://www.ema.europa.eu/en/medicines/human/EPAR/quvivig

Espie, C. A. (2022a). Standard protocol for the treatment of insomnia disorder. In C. Baglioni, C. A. Espie, & D. Riemann (Eds.), *Cognitive-behavioural therapy for insomnia (CBT-I) across the lifespan. Guidelines and clinical protocols for health professionals* (pp. 19–41). Wiley & Sons.

Espie, C. A. (2022b). The '5 principles' of good sleep health. *Journal of Sleep Research*, *31*(3), e13502. https://doi.org/10.1111/jsr.13502

Espie, C. A. (2023). Revisiting the psychological inhibition model: A conceptual framework for understanding and treating insomnia using cognitive and behavioral therapeutics (CBTx). *Journal of Sleep Research*, *2023*, e13841. https://doi.org/10.1111/jsr.13841

Espie, C. A., Emsley, R., Kyle, S. D., Gordon, C., Drake, C. L., Siriwardena, A. N., Cape, J., Ong, J. C., Sheaves, B., Foster, R., Freeman, D., Costa-Font, J., Marsden, A., & Luik, A. I. (2019). Effect of digital cognitive behavioral therapy for insomnia on health, psychological well-being, and sleep-related quality of life: A randomized clinical trial. *JAMA Psychiatry*, *76*(1), 21–30. https://doi.org/10.1001/jamapsychiatry.2018.2745

Espie, C. A., Farias Machado, P., Carl, J. R., Kyle, S. D., Cape, J., Siriwardena, A. N., & Luik, A. I. (2018). The sleep condition indicator: Reference values derived from a sample of 200 000 adults. *Journal of Sleep Research*, *27*(3), e12643. https://doi.org/10.1111/jsr.12643

Espie, C. A., Torous, J., & Brennan, T.A. (2022). Digital therapeutics should be regulated with gold-standard evidence. *Health Affairs Forefront*. doi: 10.1377/forefront.20220223.739329

Espie, C. A., Kyle, S. D., Hames, P., Gardani, M., Fleming, L., & Cape, J. (2014). The sleep condition indicator: A clinical screening tool to evaluate insomnia disorder. *BMJ Open*, *4*(3), e004183. https://doi.org/10.1136/bmjopen-2013-004183

Espie, C. A., Kyle, S. D., Miller, C. B., Ong, J. C., Hames, P., & Fleming, L. (2014). Attribution, cognition and psychopathology in persistent insomnia disorder: Outcome and mediation analysis from a randomized placebo-controlled trial of online cognitive behavioral therapy. *Sleep Medicine*, *15*(8), 913–917.

Espie, C. A., Kyle, S. D., Williams, C., Ong, J. C., Douglas, N. J., Hames, P., & Brown, J. S. (2012). A randomized, placebo-controlled trial of online cognitive behavioral therapy for chronic insomnia disorder delivered via an automated media-rich web application. *Sleep*, *35*(6), 769–781.

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Espie, C. A., Lindsay, W. R., Brooks, D. N., Hood, E. M., & Turvey, T. (1989). A controlled comparative investigation of psychological treatments for chronic sleep-onset insomnia. *Behaviour Research and Therapy*, *27*(1), 79–88. https://doi.org/10.1016/0005-7967(89)90123-x

Espie, C. A., Torous, J., & Brennan, T. A. (2022). Digital therapeutics should be regulated with gold standard evidence. *Health Affairs Forefront*. https://doi.org/10.1377/forefront.20220223.739329

Everitt, H., Baldwin, D. S., Stuart, B., Lipinska, G., Mayers, A., Malizia, A. L., Manson, C. C., & Wilson, S. (2018). Antidepressants for insomnia in adults. *Cochrane Database Systematic Reviews*, *5*, D010753.

Fan, X., Su, Z., Nie, S., Yang, J., Zhang, X., Tan, D., Xie, S., Xu, Y., Zhao, Y., Feng, L., Gu, M., & Sun, X. (2020). Efficacy and safety of chinese herbal medicine long Dan Xie Gan Tang in insomnia: A systematic review and meta-analysis. *Medicine (Baltimore)*, *99*(11), e19410.

Felder, J. N., Epel, E. S., Neuhaus, J., & Krystal, A. D. (2020 May 1). & Prather, A-A. (2020). Efficacy of digital cognitive behavioral therapy for the treatment of insomnia symptoms among pregnant women: A randomized clinical trial. *JAMA Psychiatry*, *77*(5), 484–492. https://doi.org/10.1001/jamapsychiatry.2019.4491

Feng, F., Zhang, Y., Hou, J., Cai, J., Jiang, Q., Li, X., Zhao, Q., & Li, B. A. (2018). Can music improve sleep quality in adults with primary insomnia? A systematic review and network meta-analysis. *International Journal of Nursing Studies*, *77*, 189–196. https://doi.org/10.1016/j.ijnurstu.2017.10.011

Feng, G., Han, M., Li, X., Geng, L., & Miao, Y. (2020). The clinical effectiveness of cognitive behavioral therapy for patients with insomnia and depression: A systematic review and meta-analysis. *Evid Based Complementary Alternative Medicine*, *2020*, 8071821.

Fernandez, F., & Perlis, M. L. (2023). Animal models of human insomnia. *Journal of Sleep Research*, *32*(6), e13845. https://doi.org/10.1111/jsr.13845

Fernández-San-Martín, M. I., Masa-Font, R., Palacios-Soler, L., Sancho-Gómez, P., Calbó-Caldentey, C., & Flores-Mateo, G. (2010). Effectiveness of valerian on insomnia: A meta-analysis of randomized placebo-controlled trials. *Sleep Medicine*, *11*, 505–511.

Ferracioli-Oda, E., Qawasmi, A., & Bloch, M. H. (2013). Meta-analysis: Melatonin for the treatment of primary sleep disorders. *PLoS One*, *8*, e63773.

Ford, M. E., Geurtsen, G. J., Groet, E., Rambaran Mishre, R., Bennekom, C. A. M., & Van Someren, E. J. W. (2023). A blended eHealth intervention for insomnia following acquired brain injury: A randomized controlled trial. *Journal of Sleep Research*, *32*, e13629.

Forma, F., Pratiwadi, R., El-Moustaid, F., Smith, N., Thorndike, F., & Velez, F. (2022). Network meta-analysis comparing the effectiveness of a prescription digital therapeutic for chronic insomnia to medications and face-to-face cognitive behavioral therapy in adults. *Current Medical Research Opinion*, *38*, 1727–1738.

Freeman, D., Sheaves, B., Goodwin, G. M., Yu, L. M., Nickless, A., Harrison, P. J., Emsley, R., Luik, A. I., Foster, R. G., Wadekar, V., Hinds, C., Gumley, A., Jones, R., Lightman, S., Jones, S., Bentall, R., Kinderman, P., Rowse, G., Brugha, T., … Espie, C. A. (2017). The effects of improving sleep on mental health (OASIS): A randomised controlled trial with mediation analysis. *Lancet Psychiatry*, *4*(10), 749–758.

Gao, Y., Ge, L., Liu, M., Niu, M., Chen, Y., Sun, Y., Chen, J., Yao, L., Wang, Q., Li, Z., Xu, J., Li, M., Hou, L., Shi, J., Yang, K., Cai, Y., Li, L., Zhang, J., & Tian, J. (2022). Comparative efficacy and acceptability of cognitive behavioral therapy delivery formats for insomnia in adults: A systematic review and network meta-analysis. *Sleep Medicine Reviews*, *64*, 101648. https://doi.org/10.1016/j.smrv.2022.101648

Gao, Y., Liu, M., Yao, L., Yang, Z., Chen, Y., Niu, M., Sun, Y., Chen, J., Hou, L., Sun, F., Wu, S., Zhang, Z., Zhang, J., Li, L., Li, J., Zhao, Y., Fan, J., Ge, L., & Tian, J. (2022). Cognitive behavior therapy for insomnia in cancer patients: A systematic review and network meta-analysis. *Journal of Evidence Based Medicine*, *15*, 216–229.

Garcia-Borreguero, D., Silber, M. H., Winkelman, J., Högl, B., Bainbridge, J., Buchfüherer, M., Hadjigeorgiou, G., Inoue, Y., Manconi, M., Oertel, W., Ondo, W., Winlkelmann, K., & Allen, R. P. (2016). Guidelines for the first-line treatment of restless legs syndrome/Willis-Ekbom disease, prevention and treatment of dopaminergic augmentation: A ccombined task force of the IRLSSG, EURLSSG, and the restless legs foundation. *Sleep Medicine*, *21*, 1–11.

Geiger-Brown, J. M., Rogers, V. E., Liu, W., Ludeman, E. M., Downton, K. D., & Diaz-Abad, M. (2015). Cognitive behavioral therapy in persons with comorbid insomnia: A meta-analysis. *Sleep Medicine Reviews*, *23*, 54–67.

Glass, J., Lanctôt, K. L., Herrmann, N., Sproule, B. A., & Busto, U. E. (2005). Sedative hypnotics in older people with insomnia: Meta-analysis of risks and benefits. *British Medical Journal*, *331*, 1169.

Gould, R. L., Coulson, M. C., Patel, N., Highton-Williamson, H., & Howardet, R. J. (2014). Interventions for reducing benzodiazepine use in older people: Meta-analysis of randomised controlled trials. *British Journal of Psychiatry*, *204*, 98–107.

Han, D., Cheng, J., Qu, J., Wen, X., Liu, X., Chen, Y., Wen, Y., Liu, Z., Liu, H., & Huang, Y. (2022). Effectiveness of Taijiquan in treating insomnia: A systematic review and meta-analysis of randomized controlled studies. *Frontiers in Psychiatry*, *13*, 892453. https://doi.org/10.3389/fpsyt.2022.892453 eCollection 2022.

Hasan, F., Lee, H. C., Chen, P. Y., Wang, Y. H., Yuliana, L. T., Romadlon, D. S., Tu, Y. K., & Chiu, H. Y. (2023). Comparative efficacy of hypnotics in young and middle-aged adults with insomnia: A systematic review and network meta-analysis. *Sleep Breath*, 1–10. https://doi.org/10.1007/s11325-023-02812-5

Hasan, F., Tu, Y. K., Yang, C. M., Gordon, C. J., Wu, D., Lee, H. C., Yuliana, L. T., Herawati, L., Chen, T. J., & Chiu, H. Y. (2022). Comparative efficacy of digital cognitive behavioral therapy for insomnia: A systematic review and network meta-analysis. *Sleep Medicine Reviews*, *61*, 101567.

Hauri, P. J. (1991). *Case studies in insomnia*. Plenum Medical Book Company.

Hertenstein, E., Feige, B., Gmeiner, T., Kienzler, C., Spiegelhalder, K., Johann, A., Jansson-Fröjmark, M., Palagini, L., Rücker, G., Riemann, D., & Baglioni, C. (2019). Insomnia as a predictor of mental disorders: A systematic review and meta-analysis. *Sleep Medicine Reviews*, *43*, 96–105. https://doi.org/10.1016/j.smrv.2018.10.006

Hertenstein, E., Trinca, E., Wunderlin, M., Schneider, C. L., Züst, M. A., Fehér, K. D., Su, T., van Straten, A. V., Berger, T., Baglioni, C., Johann, A., Spiegelhalder, K., Riemann, D., Feige, B., & Nissen, C. (2022). Cognitive behavioral therapy for insomnia in patients with mental disorders and comorbid insomnia: A systematic review and meta-analysis. *Sleep Medicine Reviews*, *62*, 101597. https://doi.org/10.1016/j.smrv.2022.101597

Hieu, T. H., Dibas, M., Surya, K. A., Sherif, N. A., Hashmi, M. U., Mahmoud, M., Trang, N. T. T., Abdullah, L., Nghia, T. L. B., Hirayama, K., & Huy, N. T. (2019). Therapeutic efficacy and safety of chamomile for state anxiety, generalized anxiety disorder, insomnia, and sleep quality: A systematic review and meta-analysis of randomized trials and quasi-randomized trials. *Phytotherapy Research*, *33*, 1604–1615.

Hinterberger, A., Eigl, E.-S., Hauser, T., Topalidis, P., & Schabus, M. (2022). NUKKUAA: Insomnia treatment using a novel smartphone app – First results and clinical data. In *Poster presented at the 26th congress of the European Sleep Research Society (ESRS)* September 27th – 30th 2022.

Hinterberger, A., Eigl, E.-S., Hauser, T., Topalidis, P., & Schabus, M. (2023). Evaluierung der klinischen Relevanz von NUKKUAA – einer innovativen Smartphone-App zur Behandlung von Insomnie. In *31st congress of the Austrian sleep research association*.

Ho, F. Y., Chan, C. S., Lo, W. Y., & Leung, J. C. (2020). The effect of self-help cognitive behavioral therapy for insomnia on depressive

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

symptoms: An updated meta-analysis of randomized controlled trials. *Journal of Affective Disorders*, *265*, 287–304.

Ho, F. Y., Chan, C. S., & Tang, K. N. (2016). Cognitive-behavioral therapy for sleep disturbances in treating posttraumatic stress disorder symptoms: A meta-analysis of randomized controlled trials. *Clinical Psychology Reviews*, *43*, 90–102.

Ho, F. Y., Chung, K. F., Yeung, W. F., Ng, T. H., Kwan, K. S., Yung, K. P., & Cheng, S. K. (2015). Self-help cognitive-behavioral therapy for insomnia: A meta-analysis of randomized controlled trials. *Sleep Medicine Reviews*, *19*, 17–28.

Holbrook, A. M., Crowther, R., Lotter, A., Cheng, C., & King, D. (2000). Meta-analysis of benzodiazepine use in the treatment of insomnia. *CMAJ*, *162*, 225–233.

Hu, J., Teng, J., Wang, W., Yang, N., Tian, H., Zhang, W., Peng, X., & Zhang, J. (2021). Clinical efficacy and safety of traditional Chinese medicine Xiao Yao San in insomnia combined with anxiety. *Medicine (Baltimore)*, *100*(43), e27608.

Hu, N., Ma, Y., He, J., Zhu, L., & Cao, S. (2020). Alcohol consumption and incidence of sleep disorder: A systematic review and meta-analysis of cohort studies. *Drug and Alcohol Dependence*, *217*, 108259. https://doi.org/10.1016/j.drugalcdep.2020.108259

Huang, K., Li, S., He, R., Zhong, T., Yang, H., Chen, L., Gao, H., & Jia, Y. (2022). Efficacy of cognitive behavioral therapy for insomnia (CBT-I) in older adults with insomnia: A systematic review and meta-analysis. *Australasian Psychiatry*, *30*, 592–597.

Huang, S., Huang, Q., Zhou, Z., Zhang, J., Zhan, Y., & Liang, Z. (2022). The efficacy of V. odorata extract in the treatment of insomnia: A systematic review and meta-analysis. *Frontiers in Neurology*, *13*, 730311. https://doi.org/10.3389/fneur.2022.730311

Huedo-Medina, T. B., Kirsch, I., Middlemass, J., Klonizakis, M., & Siriwardena, A. N. (2012). Effectiveness of non-benzodiazepine hypnotics in treatment of adult insomnia: Meta-analysis of data submitted to the Food and Drug Administration. *British Medical Journal*, *345*, e8343. https://doi.org/10.1136/bmj.e8343

Hürlimann, P., Bodenmann, G., Riemann, D., & Weitkamp, K. (2023). Cognitive-behavioral therapy to te treat stress and insomnia: A randomized wait list-controlled trial of two online courses. *Journal of Sleep Research*, *32*(4), e13874. https://doi.org/10.1111/jsr.13874

IQWIQ. (2021a). IQWIQ-Berichte – Nr. 1224: Kognitive Verhaltenstherapie für Insomnie – Evidenzbericht. Hrsg.: Institut für Qualität und Wortschaftlichkeit im Gesundheitswesen, Köln. https://www.iqwig.de/download/v20-07b_kognitive-verhaltenstherapie-fuer-insomnie_evidenzbericht_v1-0.pdf

IQWIQ. (2021b). IQWIQ-Berichte – Nr. 1223: Internet- oder mobilbasierte kognitive Verhaltenstherapie für Insomnie – Evidenzbericht. Hrsg.: Institut für Qualität und Wortschaftlichkeit im Gesundheitswesen, Köln. https://www.iqwig.de/download/v20-07a_internet-oder-mobilbasierte-kognitive-verhaltenstherapie-fuer-insomnie_evidenzbericht_v1-0.pdf

Irwin, M. R., Cole, J. C., & Nicassio, P. M. (2006). Comparative meta-analysis of behavioral interventions for insomnia and their efficacy in middle-aged adults an in older adults 55+ years of age. *Health Psychology*, *25*, 3–14.

Jansson-Fröjmark, M., Alfonsson, S., Bohman, B., Rozental, A., & Norell-Clarke, A. (2022). Paradoxical intention for insomnia: A systematic review and meta-analysis. *Journal of Sleep Research*, *31*, e13464.

Jernelöv, S., Blom, K., Hentati Isacsson, N., Bjurner, P., Rosén, A., Kraepelien, M., Forsell, E., & Kaldo, V. (2022). Very long-term outcome of cognitive behavioral therapy for insomnia: One- and ten-year follow-up of a randomized controlled trial. *Cognitive Behaviour Therapy*, *51*(1), 72–88.

Jespersen, K. V., Koenig, J., Jennum, P., & Vuust, P. (2015). Music for insomnia in adults. *Cochrane Database Systematic Reviews*, *8*, CD010459.

Ji, X., Ivers, H., Beaulieu-Bonneau, S., & Morin, C. M. (2021). Complementary and alternative treatments for insomnia/insomnia-depression-anxiety symptom cluster: Meta-analysis of English and Chinese literature. *Sleep Medicine Reviews*, *58*, 101445. https://doi.org/10.1016/j.smrv.2021.10144558

Johnson, J. A., Rash, J. A., Campbell, T. S., Savard, J., Gehrman, P. R., Erlis, M. L., Carlson, L. E., & Garland, S. N. (2016). A systematic review and meta-analysis of randomized controlled trials of cognitive behavior therapy for insomnia (CBT-I) in cancer survivors. *Sleep Medicine Reviews*, *27*, 20–28.

Kaldo, V., Jernelov, S., Blom, K., Ljótsson, B., Brodin, M., & Jorgensen, B. (2015). Guided internet cognitive behavioral therapy for insomnia compared to a control treatment – a randomized trial. *Behaviour Research and Therapy*, *71*, 90–100.

Khadivzadeh, T., Abdolahian, S., Ghazanfarpour, M., Kargarfard, L., Dizavandi, F. R., & Khorsand, I. (2018). A systematic review and meta-analysis on the effect of herbal medicine to manage sleep dysfunction in peri- and post-menopause. *Journal of Menopausal Medicine*, *24*, 92–99.

Kim, S. H., & Lim, J. H. (2022). Traditional east Asian herbal medicine for post-stroke insomnia: A systematic review and meta-analysis of randomized controlled trials. *International Journal of Environmental Research and Public Health*, *19*(3), 1754. https://doi.org/10.3390/ijerph19031754

Koffel, E. A., Koffel, J. B., & Gehrman, P. R. (2015). A meta-analysis of group cognitive behavioral therapy for insomnia. *Sleep Medicine Reviews*, *1*, 6–16.

Kolla, B. P., Mansukhani, S., & Mansukhani, M. P. (2016). Consumer sleep tracking devices: A review of mechanisms, validity and utility. *Expert Review Medical Devices*, *13*, 497–506.

Kraepelien, M., Blom, K., Forsell, E., Isacsson, N. H., Bjurner, P., Morin, C. M., Jernelöv, S., & Kaldo, V. (2021). A very brief self-report scale for measuring insomnia severity using two items from the insomnia severity index-development and validation in a clinical population. *Sleep Medicine*, *81*, 365–374. https://doi.org/10.1016/j.sleep.2021.03.003

Krone, L. B., Feher, K., Rivero, T., & Omlin, X. (2023). Brain stimulation techniques as novel treatment options for insomnia – A systematic review. *Journal of Sleep Research*, *2023*, e13927.

Kunz, D., Dauvilliers, Y., Benes, H., Garcia-Borreguero, D., Plazzi, G., Kinter, D. S., Coloma, P., rausch, M., Sassi-Sayadi, M., & Thein, S. (2023). Long-term safety and tolerability of daridorexant in patients with insomnia disodre. *CNS Drugs*, *37*, 93–106.

Kuriyama, A., Honda, M., & Hayashino, Y. (2014). Ramelteon for the treatment of insomnia in adults: A systematic review and meta-analysis. *Sleep Medicine*, *15*, 385–392.

Kwon, C. Y., Lee, B., Cheong, M. J., Kim, T. H., Jang, B. H., Chung, S. Y., & Kim, J. W. (2021). Non-pharmacological treatment for elderly individuals with insomnia: A systematic review and network meta-analysis. *Frontiers in Psychiatry*, *11*, 608896. https://doi.org/10.3389/fpsyt.2020.608896 eCollection 2020.

Kwon, M., Wang, J., Wilding, G., Dickerson, S. S., & Dean, G. E. (2022). Brief behavioral treatment for insomnia: A meta-analysis. *Behavioral Sleep Medicine*, *20*, 674–694.

Kyle, S. D., Aquino, M. R. J., Miller, C. B., Henry, A. L., Crawford, M., Espie, C. A., & Spielman, A. J. (2015). Towards standardisation and improved understanding of sleep restriction therapy for insomnia disoder: A systematic examination of CBT-I trial content. *Sleep Medicine Reviews*, *23*, 83–88.

Kyle, S. D., Miller, C. B., Rogers, Z., Siriwardena, A. N., Macmahon, K. M., & Espie, C. A. (2014). Sleep restriction therapy for insomnia is associated with reduced objective total sleep time, increased daytime somnolence, and objectively impaired vigilance: Implications fort he clinical management of insomnia disorder. *Sleep*, *37*, 229–237.

Kyle, S. D., Siriwardena, A. N., Espie, C. A., Yang, Y., Petrou, S., Ogburn, E., Begum, N., Maurer, L. F., Robinson, B., Gardner, C., Lee, V., Armstrong, S., Pattinson, J., Mort, S., Temple, E., Harris, V., Yu, L.-Y., Bower, P., & Aveyard, P. (2023). Clinical and cost-effectiveness of nurse-delivered sleep restriction therapy for insomnia in primary care

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

(HABIT): A pragmatic, superiority, open-label, randomised controlled trial. *Lancet*, *23*, 83–88. in press.

Lacks, P., & Rotert, M. (1986). Knowledge and practice of sleep hygiene techniques in insomniacs and good sleepers. *Behavior Research and Therapy*, *24*, 365–368.

Lam, T. H., Chung, K. F., Yeung, W. F., Yu, B. Y., Yung, K. P., & Ng, T. H. (2015). Hypnotherapy for insomnia: A systematic review and meta-analysis of randomized controlled trials. *Complementary Therapy and Medicine*, *23*, 719–732.

Leach, M. J., & Page, A. T. (2015). Herbal medicine for insomnia: A systematic review and meta-analysis. *Sleep Medicine Reviews*, *24*, 1–12.

LeBlanc, E. S., Smith, N. X., Nichols, G. A., Allison, M. J., & Clarke, G. N. (2018). Insomnia is associated with an increased risk of type 2 diabetes in the clinical setting. *British Medical Journal, Open Diabetes Research & Care*, *6*(1), e000604. https://doi.org/10.1136/bmjdrc-2018-000604

Lebrun, C., Gely-Nargeot, M. C., Rossignol, A., Geny, C., & Bayard, S. (2020). Effeicacy of cognitive behavioral therapy for insomnia comorbid to Parkinsons's disease: A focus on psychological and daytime functioning with a single-case design with multiple baselines. *Journal of Clinical Psychology*, *76*, 356–376.

Leerssen, J., Lakbila-Kamal, O., Dekkers, L., Ikelaar, S. L., Albers, A. C., Blanken, T. F., Lancee, J., Van Der Lande, G. J., Maksimovic, T., Mastenbroek, S. E., & Reesen, J. E. (2022). Treating insomnia with high risk of depression using therapist-guided digital cognitive, behavioral, and circadian rhythm support interventions to prevent worsening of depressive symptoms: A randomized controlled trial. *Psychotherapy and Psychosomatics*, *91*, 168–179.

Li, F., Xu, B., Shi, H., Zhang, T., Song, Z., Chen, Y., Liu, L., & Wang, P. (2020). Efficacy and safety of TCM Yangxin Anshen therapy for insomnia: A systematic review and meta-analysis. *Medicine (Baltimore)*, *99*, e19330.

Li, H., Chen, J., Xu, G., Duan, Y., Huang, D., Tang, C., & Liu, J. (2020). The effect of tai chi for improving sleep quality: A systematic review and meta-analysis. *Journal of Affective Disorders*, *274*, 1102–1112.

Li, L., Gan, Y., Zhou, X., Jiang, H., Zhao, Y., Tian, Q., He, Y., Liu, Q., Mei, Q., Wu, C., & Lu, Z. (2021). Insomnia and the risk of hypertension: A meta-analysis of prospective cohort studies. *Sleep Medicine Reviews*, *56*, 101403. https://doi.org/10.1016/j.smrv.2020.101403

Li, S., Li, Z., Wu, Q., Liu, C., Zhou, Y., Chen, L., Lan, W., Li, Y., Han, S., & Chen, Z. (2021). Effect of exercise intervention on primary insomnia: A meta-analysis. *Journal of Sports Medicine*, *61*, 857–866.

Lian, J., Zhong, Y., Li, H., Yang, S., Wang, J., Li, X., Zhou, X., & Chen, G. (2022). Effects of saffron supplementation on improving sleep quality: A meta-analysis of randomized controlled trials. *Sleep Medicine*, *92*, 24–33.

Liang, L., Huang, Y., Xu, R., Wei, Y., Xiao, L., & Wang, G. (2019). Eszopiclone for the treatment of primary insomnia: A systematic review and meta-analysis of double-blind, randomized, placebo-controlled trials. *Sleep Medicine*, *62*, 6–13.

Lin, Y. H., Chen, C., Zhao, X., Mao, Y. F., Xiang, G. X., Yang, M. Q., & Song, Y. M. (2021). Efficacy and safety of Banxia formulae for insomnia: A systematic review and meta-analysis of high-quality randomized controlled trials. *Evidence Based Complementary and Alternative Medicine*, *833168*, 1–18.

Liu, J., & Wang, L. N. (2012). Ramelteon in the treatment of chronic insomnia: Systematic review and meta-analysis. *International Journal of Clinical Practice*, *2*(66), 867–873.

Liu, Y., Xu, X., Dong, M., Jia, S., & Wie, Y. (2017). Treatment of insomnia with tricyclic antidepressants: A meta-analysis of polysomnographic randomized controlled trials. *Sleep Medicine*, *34*, 126–133.

Lopez, R., Evangelista, E., Barateau, L., Chenini, S., Bosco, A., Billiard, M., Bonte, A. D., Béziat, S., Jaussent, I., & Dauvilliers, Y. (2019). French language cnline cognitive behavioral therapy for insomnia disorder: A randomized controlled trial. *Frontiers in Neurology*, *10*, 1273. https://doi.org/10.3389/fneur.2019.01273

Lorenz, N., Heim, E., Roetger, A., Birrer, E., & Maercker, A. (2019). Randomized controlled trial to test the efficacy of an unguided online intervention with automated feedback for the treatment of insomnia. *Behavioural and Cognitive Psychotherapy*, *47*, 287–302.

Lorenz, W., Ollenschläger, G., & Geradts, M. (2001). Das Leitlinien Manual: Entwicklung und Implementierung von Leitlinien in der Medizin. *Zeitschrift für Ärztliche Fortbildung und Qualitätssicherung im Gesundheitswesen*, *95*, 1–84.

Luan, X., Zhang, X., & Zhou, Y. (2022). The role and clinical observation of traditional Chinese medicine in relieving senile insomnia: A systematic review and meta-analysis. *Biomedical Research International*, *9484095*. https://doi.org/10.1155/2022/9484095

Luik, A. I., Machado, P. F., Siriwardena, N., & Espie, C. A. (2019). Screening for insomnia in primary care: Using a two-item version of the sleep condition indicator. *British Journal of General Practice*, *69*(679), 79–80. https://doi.org/10.3399/bjgp19X701045

Ma, Y., Hall, D. L., Ngo, L. H., Liu, Q., Bain, P. A., & Yeh, G. Y. (2021). Efficacy of cognitive behavioral therapy for insomnia in breast cancer: A meta-analysis. *Sleep Medicine Reviews*, *55*, 101376.

Marschall, J., Hildebrandt, S., Sydow, H., & Nolting, H. D. (2017). *Gesundheitsreport 2017*. Medhochzwei Verlag GmbH.

Maruani, J., Reynaud, E., Chambe, J., Palagini, L., Borigin, P., & Geoffroy, P. A. (2023). Efficacy of melatonin and ramelteon for the actute and long-term-management of insomnia disorder in adults: A systematic review and meta-analysis. *Journal of Sleep Research*, *2023*, 23–195. acc.

Marupuru, S., Arku, D., Campbell, A. M., Slack, M. K., & Lee, J. K. (2022). Use of melatonin and/or ramelteon for the treatment of insomnia in older adults: A systematic review and meta-analysis. *Journal of Clinical Medicine*, *11*, 5138.

Maurer, L., Espie, C. A., & Kyle, S. D. (2018). How does sleep restriction therapy for insomnia work? A systematic review of mechanistic evidence and the introduction of the triple-R model. *Sleep Medicine Reviews*, *41*, 127–138.

Maurer, L., Espie, C. A., Omlin, X., Emsley, R., & Kyle, S. D. (2022). The effect of sleep restriction therapy for insomnia on sleep pressure and arousal: A randomized controlled mechanistic trial. *Sleep*, *45*, zsab223.

Maurer, L. F., Espie, C. A., Omlin, X., Reidse, M. J., Sharman, R., Gavriloff, D., Emsley, R., & kyle, S. D. (2020). Isolating the role of time in bed restriction in the treatment of insomnia: A randomized, controlled, dismantling trial comparing sleep restriction therapy with time in bed regularization. *Sleep*, *43*, zsaa096.

Maurer, L. F., Schneider, J., Miller, C. B., Espie, C. A., & Kyle, S. D. (2021). The clinical effects of sleep restriction therapy for insomnia: A meta-analysis of randomised controlled trials. *Sleep Medicine Reviews*, *58*, 101493.

McCleery, J., Cohen, D. A., & Sharpley, A. L. (2014). Pharmacotherapies for sleep disturbances in Alzheimer's disease. *Cochrane Database Systematic Reviews*, *3*, CD009178.

Mercier, J., Savard, J., & Bernard, P. (2017). Exercise interventions to improve sleep in cancer patients: A systematic review and meta-analysis. *Sleep Medicine Reviews*, *36*, 43–56.

Mignot, E., Mayleben, D., Fietze, I., Leger, D., Zammit, G., Bassetti, C. L. A., Pain, S., Kinter, D. S., & Roth, T. (2022). Safety and efficacy of daridorexant in patients with insomnia disorder: Results from two multicenter, randomised, double-blind, placebo-controlled, phase 3 trials. *Lancet Neurology*, *21*, 125–139.

Miller, C. B., Espie, C. A., Epstein, D. R., Friedman, L., Morin, C. M., Pigeon, W. R., Spielman, A. J., & Kyle, S. D. (2014). The evidence base of sleep restriction therapy for treating insomnia disorder. *Sleep Medicine Reviews*, *18*, 415–424.

Mitchell, L. J., Bisdounis, L., Ballesio, A., Omlin, X., & Kyle, S. D. (2019). The impact of cognitive behavioural therapy for insomnia on objective sleep parameters: A meta-analysis and systematic review. *Sleep Medicine Reviews*, *47*, 90–102.

Montgomery, P., & Dennis, J. (2004). A systematic review of non-pharmacological therapies for sleep problems in later life. *Sleep Medicine Reviews*, *8*, 47–62.

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Morin, C. M., Belleville, G., Belanger, L., & Ivers, H. (2011). The insomnia severity index: Psychometric indicators to detect insomnia cases and evaluate treatment response. *Sleep*, *34*, 601–608.

Morin, C. M., Colecchi, C., Stone, J., Sood, R., & Brink, D. (1999). Behavioral and pharmacological therapies for late-life insomnia: A randomized controlled trial. *Journal of the American Medical Association*, *281*, 991–999.

Morin, C. M., Culbert, J. P., & Schwartz, S. M. (1994). Nonpharmacological interventions for insomnia: A meta-analysis of treatment efficacy. *American Journal of Psychiatry*, *151*, 1172–1180.

Morin, C. M., Edinger, J. D., Beaulieu-Bonneau, S., Ivers, H., Krystal, A. D., Guay, B., Bélanger, L., Cartwright, A., Simmons, B., Lamy, M., & Busby, M. (2020). Effectiveness of sequential psychological and medication therapies for insomnia disorder: A randomized clinical trial. *JAMA Psychiatry*, *77*, 1107–1115.

Morin, C. M., Inoue, Y., Kushida, C., Poyares, D., & Winkelman, J. (2021). Endorsement of the european insomnia guideline for the diagnosis and treatment of insomnia by the world sleep society. *Sleep Medicine*, *81*, 124–126.

Morin, C. M., Vallières, A., Guay, B., Ivers, H., Savard, J., Mérette, C., Bastien, C., & Baillargeon, L. (2009). Cognitive behavioral therapy, singly and combined with medication, for persistent insomnia. *Journal of the American Medical Association*, *301*, 2005–2015.

Munirah, M. P., Norhayati, M. N., & Noraini, M. (2022). Crocus sativus for insomnia: A systematic review and meta-analysis. *International Journal of Environmental Research and Public Health*, *19*, 11658.

Murtagh, D. R., & Greenwood, K. M. (1995). Identifying effective psychological treatments for insomnia: A meta-analysis. *Journal of Consulting and Clinical Psychology*, *63*, 79–89.

Ni, X., Shergis, J. L., Guo, X., Zhang, A. L., Li, Y., Lu, C., & Xue, C. C. (2015). Updated clinical evidence of Chinese herbal medicine for insomnia: A systematic review and meta-analysis of randomized controlled trials. *Sleep Medicine*, *16*, 1462–1481.

Nowell, P. D., Mazumdar, S., Buysse, D. J., Dew, M. A., Reynolds, C. F., 3rd, & Kupfer, D. J. (1997). Benzodiazepines and zolpidem for chronic insomnia: A meta-analysis of treatment efficacy. *Journal of the American Medical Association*, *278*, 2170–2177.

Okajima, I., Komada, Y., & Inoue, Y. (2011). A meta-analysis on the treatment effectiveness of cognitive behavioral therapy for primary insomnia. *Sleep and Biological Rhythms*, *9*, 24–34.

Oyekan, P. J., Gorton, H. C., & Copeland, C. S. (2021). Antihistamine-related deaths in England: Are their high safety profiles of antihistamines leading to their unsafe use? *British Journal of Clinical Pharmacology*, *10*, 3978–3987. https://doi.org/10.1111/bcp.14819

Palagini, L., Geoffroy, P. A., Gemignani, A., & Riemann, D. (2023). Genetic and epigenetic potential mechanisms in insomnia: A systematic review. *Journal of Sleep Research*, *2023*, e13868. https://doi.org/10.1111/jsr.13868

Pallesen, S., Nordhus, I. H., & Kvale, G. (1998). Nonpharmacological interventions for insomnia in older adults: A meta-analysis of treatment efficacy. *Psychotherapy: Theory, Research and Practical Training*, *35*, 472–482.

Perlis, M. L., Gehrman, P., & Riemann, D. (2008). Intermittent and long-term use of sedative hypnotics. *Current Pharmaceutical Design*, *14*, 3456–3465.

Perney, P., & Lehert, P. (2018). Insomnia in alcohol-dependent patients: Prevalence, risk factors and Acamprosate effect: An individual patient data meta-analysis. *Alcohol and Alcoholism (Oxford, Oxfordshire)*, *53*(5), 611–618. https://doi.org/10.1093/alcalc/agy013

Peter, L., Reindl, R., Zauter, S., Hillemacher, T., & Richter, K. (2019). Effectiveness of an online CBT-I intervention and a face-to-face treatment for shift work sleep disorder: A comparison of sleep diary data. *International Journal of Environmental Research and Public Health*, *16*, 3081. https://doi.org/10.3390/ijerph16173081

Philip, P., Dupuy, L., Sagaspe, P., de Sevin, E., Auriacombe, M., Taillard, J., Micoulaud-Franchi, J. A., & Morin, C. M. (2022). Efficacy of a smartphone-based virtual companion to treat insomniac complaints in the general population: Sleep diary monitoring versus an internet autonomous intervention. *Journal of Clinical Medicine*, *11*(15), 4387.

Quaseem, A., Kansagara, D., Forciea, M. A., Cooke, M., & Denberg, T. D. (2016). Management of chronic insomnia disorder in adults: A clinical practice guideline from the American College of Physicians. *Annals of Internal Medicine*, *165*, 125–133.

Rapoport, M. J., Lanctot, K. L., Streiner, D. L., Bédard, M., Vingilis, E., Murray, B., Schaffer, A., Shulman, K. L., & Herrmann, N. (2009). Benzodiazepine use and driving: A meta-analysis. *Journal of Clinical Psychiatry*, *70*, 663–673.

Reffi, A. N., Kalmbach, D. A., Cheng, P., & Drake, C. L. (2023). The sleep response to stress: How sleep reactivity can help us prevent insomnia and promote resilience. *Journal of Sleep Research*, *2023*, e13892. https://doi.org/10.1111/jsr.13892

Riemann, D., Baglioni, C., Bassetti, C., Bjorvatn, B., Dolenc Groselj, L., Ellis, J., Espie, C., Garcia-Borreguero, D., Gjerstad, M., Gonçalves, M., Hertenstein, E., Jansson-Fröjmark, M., Jennum, P., Léger, D., Nissen, C., Parrino, L., Paunio, T., Pevernagie, D., Verbraecken, J., … Spiegelhalder, K. (2017). European guideline for the diagnosis and treatment of insomnia. *Journal of Sleep Research*, *26*, 675–700.

Riemann, D., Baum, E., Cohrs, S., Crönlein, T., Hajak, G., Hertenstein, E., Klose, P., Langhorst, J., Mayer, G., Nissen, C., Pollmächer, T., Rabstein, S., Schlarb, A., Sitter, H., Weess, H. G., Wetter, T., & Spiegelhalder, K. (2017). S3-Leitlinie Nicht erholsamer Schlaf/Schlafstörungen – Kapitel Insomnie bei Erwachsenen, Update 2016. *Somnologie*, *21*, 2–44.

Riemann, D., Benz, F., Blanken, T., Espie, C. A., Dressle, R., Johann, A. F., Henry, A., Kyle, S. D., Leerssen, J., Wassing, R., Spiegelhalder, K., & van Someren, E. (2022). Insomnia disorder: State of the science and challenges for the future. *Journal of Sleep Research*, *31*, e13604.

Riemann, D., & Nissen, C. (2012). Sleep and psychotropic drugs. In C. A. Espie & C. M. Morin (Eds.), (Hrsg.) *Oxford handbook of sleep and sleep disorders* (pp. 190–222). Oxford Press.

Rösler, L., van der Lane, G., Leerssen, J., Cox, R., Ramautar, J. R., & van Someren, E. J. W. (2023). Actigraphy in studies on insomnia: Worth the effort? *Journal of Sleep Research*, *32*, e13750.

Rösner, S., Englbrecht, C., Wehrle, R., Hajak, G., & Soyka, M. (2018). Eszopiclone for insomnia (reveiw). *Cochrane database of Systematic Reviews*, *2018*(10), CD010703. https://doi.org/10.1002/146511858

Rubio-Arias, J. A., Marin-Cascales, E., Ramos-Campo, D. J., Hernandez, A. V., & Perez-Lopez, F. R. (2017). Effect of exercise on sleep quality and insomnia in middle-aged women: A systematic review and meta-analysis of randomized controlled trials. *Maturitas*, *100*, 49–56.

Schabus, M., Eigl, E.-S., Hinterberger, A., Hauser, T., & Topalidis, P. (2022). Nukkuaa – A novel mobile phone application for tracking sleep and treating insomnia: A first efficacy study. In *Poster presented at the 26th conference of the European Sleep Research Society (ESRS) 2022*.

Schierenbeck, T., Riemann, D., Berger, M., & Hornyak, M. (2008). Effect of illicit recreational drugs upon sleep: Cocaine, ecstasy and marijuana. *Sleep Medicine Reviews*, *12*, 381–390.

Schneider, C., Hertenstein, E., Fehér, K. D., Maier, J. G., Cantisani, A., Moggi, F., Berger, T., & Nissen, C. (2020). Become your own SLEEPexpert: Design, implementation and first evaluation of a pragmatic behavioral treatment program for insomnia in patients with acute psychiatric disorders. *SLEEP Advances* https://academic.oup.com/sleepadvances/article/1/1/zpaa005/6016467?login=true

Scott, H., Bensen-Boakes, D. B., Lovato, N., Reynolds, A., Perlis, M., & Lack, L. (2023). The efficacy of intensive sleep retraining for insomnia: A systematic review and research agenda. *Journal of Sleep Research*, *32*, e13894. https://doi.org/10.1111/jsr.13894

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Selvanathan, J., Pham, C., Nagappa, M., Peng, P. W. H., Englesakis, M., Espie, C. A., Morin, C. M., & Chung, F. (2021). Cognitive behavioral therapy for insomnia in patients with chronic pain – A systematic review and meta-analysis of randomized controlled trials. *Sleep Medicine Reviews*, *60*, 101460.

Seyffert, M., Lagisetty, P., Landgraf, J., Chopra, V., Pfeiffer, P. N., Conte, M. L., & Rogers, M. A. (2016). Internet-delivered cognitive behavioral therapy to treat insomnia: A systematic review and meta-analysis. *PLoS One*, *11*, e0149139.

Shahid, A., Wilkinson, K., Marcu, S., & Shapiro, C. (2012). *Stop, that and one hundred other sleep scales*. Springer.

Shinjyo, N., Waddell, G., & Green, J. (2020). Valerian root in treating sleep problems and associated disorders – A systematic review and meta-analysis. *Journal of Evidence based Integrative Medicine*, *25*, 2515690X20967323.

Simon, L., Reimann, J., Steubl, L. S., Stach, M., Spiegelhalder, K., Sander, L. B., Baumeister, H., Messner, E. M., & Terhorst, Y. (2023). Help for insomnia from the appstore? A standardized rating of mobile health applications claiming to target insomnia. *Journal of Sleep Research*, *32*, e13642.

Simon, L., Steinmetz, L., Feige, B., Benz, F., Spiegelehalder, K., & Baumeister, H. (2023). Comparative efficacy of onsite, digital and other settings for cognitive behavioral therapy for insomnia: A systematic review and network meta-analysis. *Scientific Reports*, *13*, 1929.

Smith, M. T., McCrae, C. S., Cheung, J., Martin, J. L., Harrod, C. G., Heald, J. L., & Carden, K. A. (2018). Use of actigraphy for the evaluation of sleep disorders and circadian rhythm sleep-wake disorders: An American Academy of sleep medicine systematic review, meta-analysis, and GRADE assessment. *Journal of Clinical Sleep Medicine*, *14*, 1209–1230.

Sofi, F., Cesari, F., Casini, A., Macchi, C., Abbate, R., & Gensini, G. F. (2014). Insomnia and risk of cardiovascular disease: A meta-analysis. *European Journal of Preventive Cardiology*, *21*(1), 57–64. https://doi.org/10.1177/2047487312460020

Soh, H. L., Ho, R. C., Ho, C. S., & Tam, W. W. (2020). Efficacy of digital cognitive behavioural therapy for insomnia: A meta-analysis of ransomised controlled trials. *Sleep Medicine*, *75*, 315–325.

Soldatos, C. R., Dikeos, D. G., & Whitehead, A. (1999). Tolerance and rebound insomnia with rapidly eliminated hypnotics: A meta-analysis of sleep laboratory studies. *International Clinical Psychopharmacology*, *14*(5), 287–303.

Soyka, M. (2021). Missbrauch und Abhängigkeit von Benzodiazepinen und Z-Drugs. *Nervenheilkunde*, *40*, 636–646.

Spiegelhalder, K., Baum, E., Becker, M., Cornaro, C., Crönlein, T., Frase, L., Harth, V., Hertenstein, E., Johann, A., Klose, P., Mertel, I., Kunz, D., Langhorst, J., Maurer, J. T., Mayer, G., Nissen, C., Pietrowsky, A., Pollmächer, T., Schumacher, C., … Riemann, D. (2023). *Leitlinie "Insomnie bei Erwachsenen" – Update 2022. (AWMF-Registernummer 063–003). Somnologie in Press*.

Spielman, A. J., Caruso, L. S., & Glovinsky, P. A. (1987). A behavioral perspective on insomnia treatment. *Psychiatric Clinics of North America*, *10*, 541–553.

Squires, L. R., Rash, J. A., Fawcett, J., & Farland, S. N. (2022). Systematic review and meta-analysis of cognitive-behavioural therapy for insomnia on subjective and actigraphy-measured sleep and comorbid symptoms in cancer survivors. *Sleep Medicine Reviews*, *63*, 101615.

Steinmetz, L., Simon, L., Feige, B., Riemann, D., Johann, A. F., Ebert, D. D., Baumeister, H., Benz, F., & Spiegelhalder, K. (2023). Network meta-analysis examining efficacy of components of cognitive behavioural therapy for insomnia. Clinical Psychology Reviews, submitted.

Steinmetz, L., Simon, L., Riemann, D., Feige, B., Johann, A. F., Baumeister, H., Ebert, D., Benz, F., & Spiegelhalder, K. (2022). Network meta-analysis examining the efficacy of components of cognitive behavioural therapy for insomnia. *Journal of Sleep Research*, *31*(S1). https://doi.org/10.1111/jsr.13740

Stenberg, J.-H., Ritola, V., Joffe, G., Saarni, S., & Rosenström, T. (2022). Effectiveness of mobile-delivered, therapist-assisted cognitive behavioral therapy for insomnia in nationwide routine clinical care in Finland. *Journal of Clinical Sleep Medicine*, *18*(11), 2643–2651.

Stranks, E. K., & Crowe, S. F. (2014). The acute cognitive effects of zopiclone, zolpidem, zaleplon, and eszopiclone: A systematic review and meta-analysis. *Journal of Clinical and Experimental Neuropsyvchology*, *36*, 691–700.

Ström, L., Pettersson, R., & Andersson, G. (2004). Internet-based treatment for insomnia: A controlled evaluation. *Journal of Consulting and Clinical Psychology*, *72*, 113–120.

Sunnhed, R., Hesser, H., Andersson, G., Carlbring, P., Morin, C. M., Harvey, A., & Jansoon-Fröjmark, M. (2020). Comparing internet-delivered cognitive therapy and behavior therapy with telephone support for insomnia disorder: A randomized controlled trial. *Sleep*, *43*, zsz245.

Sweetman, A., Farrell, S., Wallace, D., & Crawford, M. (2023). The effect of cognitive behavioural therapy for insomnia in people with co-morbid insomnia and sleep apnoe: A systematic review and meta-analysis. *Journal of Sleep Research*, *32*, e13847. https://doi.org/10.1111/jsr.13847

Sweetman, A., Lack, L., McEvoy, R. D., Smith, S., Eckert, D. J., Osman, A., varberry, J. C., Wallace, D., Nguyen, P. D., & Catcheside, P. (2021). Bi-directional relationships between co-morbid insomnia and sleep apnea (COMISA). *Sleep Medicine Reviews*, *60*, 101519. https://doi.org/10.1016/j.smrv.2021.101519

Tang, N., Saconi, B., Jansson-Fromark, M., Ong, J., & Carney, C. (2023). Cognitive factors and processes in models of insomnia: A systematic review. *Journal of Sleep Research*, *32*, e13923.

Tang, N. K., & Harvey, A. G. (2006). Altering misperception of sleep in insomnia: Behavioral experiment versus verbal feedback. *Journal of Consulting and Clinical Psychology*, *74*, 767–776.

Tang, N. K., Lereya, S. T., Boulton, H., Miller, M. A., Wolke, D., & Cappuccio, F. P. (2015). Nonpharmacological treatments of insomnia for long-term painful conditions: A systematic review and meta-analysis of patient-reported outcomes in randomized controlled trials. *Sleep*, *38*, 1751–1764.

Thakral, M., von Korff, M., McCurry, S. M., Morin, C. M., & Vitiello, M. V. (2020). Changes in dysfunctional beliefs about sleep after cognitive behavioral therapy for insomnia: A systematic literature review and meta-analysis. *Sleep Medicine Reviews*, *49*, 101230.

Thiart, H., Lehr, D., Ebert, D. D., Berking, M., & Riper, H. (2015). Log in and breathe out: Internet-based recovery training for sleepless employees with work-related strain – results of a randomized controlled trial. *Scandinavian Journal of Work, Environment & Health*, *41*(2), 164–174. https://doi.org/10.5271/sjweh.3478

Thorndike, F. P., Ritterband, L. M., Gonder-Frederick, L. A., Lord, H. R., Ingersoll, K. S., & Morin, C. M. (2013). A randomized controlled trial of an internet intervention for adults with insomnia: Effects on comorbid psychological and fatigue symptoms. *Journal of Clinical Psychology*, *69*(10), 1078–1093. https://doi.org/10.1002/jclp.22032

Trauer, J. M., Qian, M. Y., Doyle, J. S., Rajaratnam, S. M., & Cunnington, D. (2015). Cognitive behavioral therapy for chronic insomnia: A systematic review and meta-analysis. *Annals of Internal Medicine*, *163*, 191–204.

Treves, N., Perlman, A., Kolenberg, G. L., Asaly, A., & Matok, I. (2018). Z-drugs and risk for falls and fractures in older adults – A systematic review and meta-analysis. *Age and Ageing*, *47*, 201–208.

Turner, R. M., & Ascher, L. M. (1979). Controlled comparison of progressive relaxation, stimulus control, and paradoxical intention therapies for insomnia. *Journal of Consulting and Clinical Psychology*, *47*(3), 500–508. https://doi.org/10.1037//0022-006x.47.3.500

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

van der Zweerde, T., Bisdounis, L., Kyle, S. D., Lancee, J., & van Straten, A. (2019). Cognitive behavioral therapy for insomnia: A meta-analysis of long-term effects in controlled studies. *Sleep Medicine Reviews*, *48*, 101208.

Van der Zweerde, T., Lancee, J., Slottje, P., Bosmans, J. E., Van Someren, E. J. W., & van Straten, A. (2020). Nurse-guided internet-delivered cognitive behavioral therapy for insomnia in general practice: Results from a pragmatic randomized clinical trial. *Psychotherapy and Psychosomatics*, *89*(3), 174–184.

Van der Zweerde, T., van Straten, A., Effting, M., Kyle, S. D., & Lancee, J. (2019). Does online insomnia treatment reduce depressive symptoms? A randomized controlled trial in individuals with both insomnia and depressive symptoms. *Psychological Medicine*, *49*(3), 501–509.

Van Maanen, A., Meijer, A. M., van der Heijden, K. B., & Oort, F. J. (2016). The effects of light therapy on sleep problems: A systematic review and meta-analysis. *Sleep Medicine Reviews*, *29*, 52–62.

Van Someren, E. J. W. (2021). Brain mechanisms of insomnia: New perspectives on causes and consequences. *Physiological Reviews*, *101*(3), 995–1046. https://doi.org/10.1152/physrev.00046.2019

Van Straten, A., & Cuijpers, P. (2009). Self-help therapy for insomnia: A meta-analysis. *Sleep Medicine Reviews*, *13*, 61–71.

Van Straten, A., Emmelkamp, J., de Wit, J., Lancee, J., Andersson, G., van Someren, E. W. J., & Cuijpers, P. (2014). Guided internet-delivered cognitive behavioral treatment for insomnia: A randomized trial. *Psychological Medicine*, *44*(7), 1521–1532.

Van Straten, A., van der Zweerde, T., Kleiboer, A., Cuijpers, P., Morin, C. M., & Lancee, J. (2018). Cognitive and behavioral therapies in the treatment of insomnia: A meta-analysis. *Sleep Medicine Reviews*, *38*, 3–16.

Vande Griend, J. P., & Anderson, S. L. (2012). Histamine-1 receptor antagonism for treatment of insomnia. *Journal of the American Pharmacological Association*, *52*, e210–e219.

Wang, C., Yang, Y., Ding, X., Li, J., Zhou, X., Teng, J., & Qi, X. (2021). Efficacy and safety of Shumian capsules in treating insomnia: A systematic review and meta-analysis. *Medicine (Baltimore)*, *100*, e28194.

Wang, X., Ju, J., Li, J., Fan, Y., & Xu, H. (2020). Chaihu Longgu Muli decoction, a Chinese herbal formula, for the treatment of insomnia: A systematic review and meta-analysis. *Medicine (Baltimore)*, *99*, e22462.

Wang, Y. Y., Wang, F., Zheng, W., Zhang, L., Ng, C. H., Ungvari, G. S., & Xiang, Y. T. (2020). Mindfulness-based interventions for insomnia: A meta-analysis of randomized controlled trials. *Behavioiral Sleep Medicine*, *18*, 1–9.

Wilson, S., Anderson, K., Baldwin, D., Dijk, D.-J., Espie, A., Espie, C., & Sharpley, A. (2019). British Association for Psychopharmacology consensus statement on evidence-based treatment of insomnia, parasomnias and circadian rhythm disorders: An update. *Journal of Psychopharmacology*, *33*(8), 923–947. https://doi.org/10.1177/0269881119855343

Winkler, A., Auer, C., Doering, B. K., & Rief, W. (2014). Drug treatment of primary insomnia: A meta-analysis of polysomnographic randomized controlled trials. *CNS Drugs*, *28*, 799–816.

World Health Organization (WHO). International classification of diseases, 10th edition. (1994). *In: World Health Organization (WHO) (editor)*. World Health Organization.

World Health Organization (WHO). International classification of diseases, 11th edition. (2022). *In: World Health Organization (WHO) (editor)*. World Health Organization.

Wu, J. Q., Appleman, E. R., Salazar, R. D., & Ong, J. C. (2015). Cognitive behavioral therapy for insomnia comorbid with psychiatric and medical conditions. *JAMA Internal Medicine*, *175*, 1461–1472.

Xiang, T., Cai, Y., Hong, Z., & Pan, J. (2021). Efficacy and safety of zolpidem in the treatment of insomnia disorder for one month: A meta-analysis of a randomized controlled trial. *Sleep Medicine*, *87*, 250–256.

Xu, D., Cardell, E., Broadley, S. A., & Sun, J. (2021). Efficacy of face-to-face delivered cognitive behavioral therapy in improving health status of patients with insomnia: A meta-analysis. *Frontiers in Psychiatry*, *12*, 798453.

Yang, X. Q., Liu, L., Ming, S. P., Fang, J., & Wu, D. N. (2019). Tian Wang Bi Xin Dan for insomnia: A systematic review of efficacy and safety. *Evidence Based Complement Ary and Alternative Medicine*, *2019*, 4260801.

Ye, Y. Y., Zhang, Y. F., Chen, J., Liu, J., Li, X. J., Liu, Y. Z., Lang, Y., Lin, L., Yang, X. J., & Jiang, X. J. (2015). Internet-based cognitive behavioral therapy for insomnia (ICBT-I) improves comorbid anxiety and depression – a meta-analysis of randomized controlled trials. *PLoS One*, *10*, e0142258.

Yeung, V., Sharpe, L., Glozier, N., Hackett, M. L., & Colagiuri, B. (2018). A systematic review and meta-analysis of placebo versus no treatment for insomnia symptoms. *Sleep Medicine Reviews*, *38*, 17–27.

Yeung, W. F., Chung, K. F., Yung, K. P., & Ng, T. H. Y. (2015). Doxepin for insomnia: A systematic review of randomized placebo-controlled trials. *Sleep Medicine Reviews*, *19*, 75–83.

Yi, X. Y., Ni, S. F., Ghadami, M. R., Meng, H. Q., Chen, M. Y., Kuang, L., Zhang, Y. Q., Zhang, L., & Zhou, X. Y. (2018). Trazodone for the treatment of insomnia: A meta-analysis of randomized placebo-controlled trials. *Sleep Medicine*, *45*, 25–32.

Yoon, J. H., Kim, E. H., Park, S. B., Lee, J. Y., & Yoon, S. W. (2021). Traditional herbal medicine for insomnia in patients with cancer: A systematic review and meta-analysis. *Frontiers in Pharmacology*, *12*, 753140.

Yu, H., Zhang, Y., Liu, Q., & Yan, R. (2021). Efficacy of online and face-to-face cognitive behavioral therapy in the treatment of neurological insomnia: A systematic review and meta-analysis. *Annals of Palliative Medicine*, *10*, 10684–10696.

Yue, J. L., Chang, X. W., Zheng, J. W., Shi, L., Xiang, Y. J., Que, J. Q., Yuan, K., Deng, J. H., Teng, T., Li, Y. Y., Sun, W., Sun, H. Q., Vitiello, M., Tang, X. D., Zhou, X. Y., Bao, Y. P., Shi, J., & Lu, L. (2023). Efficacy and tolerability of pharmacological treatments for insomnia in adults: A systematic review and network meta-analysis. *Sleep Medicine Reviews*, *68*, 101746.

Zachariae, R., Lyby, M. S., Ritterband, L. M., & O'Toole, M. S. (2016). Efficacy of internet-delivered cognitive-behavioral therapy for insomnia – A systematic review and meta-analysis of randomized controlled trials. *Sleep Medicine Reviews*, *30*, 1–10.

Zakiei, A., Khazaie, H., Rostampour, M., Lemola, S., Esmaeili, M., Dürsteler, K., Brühl, A. B., Sadeghi-Bahmani, D., & Brand, S. (2021). Acceptance and commitment therapy (ACT) improves sleep quality, experiential avoidance, and emotion regulation in individuals with insomnia – Results from a randomized interventional study. *Life*, *11*, 133.

Zhang, H., Liu, P., Wu, X., Zhang, Y., & Cong, D. (2019). Effectiveness of Chinese herbal medicine for patients with primary insomnia: A PRISMA-compliant meta-analysis. *Medicine (Baltimore)*, *98*, e15967.

Zhang, W., Chen, X. Y., Su, S. W., Jia, Q. Z., Ding, T., Zhu, Z. N., & Zhang, T. (2016). Exogenous melatonin for sleep disorders in neurodegenerative diseases: A meta-analysis of randomized clinical trials. *Neurological Science*, *37*, 57–65.

Zhang, Y., Ren, R., Yang, L., Zhang, H., Shi, Y., Shi, J., Sanford, L. D., Lu, L., Vitiello, M. V., & Tang, X. (2022). Comparative efficacy and acceptability of psychotherapies, pharmacotherapies, and their combination for the treatment of adult insomnia: A systematic review and network meta-analysis. *Sleep Medicine Reviews*, *65*, 101687.

Zhang, Y., ren, R., Zhou, J., Zhang, J., Wing, Y. K., Sanford, L., & Tang, X. (2019). Worldwide and regional prevalence rates of co-occurence of insomnia and insomnia symptoms with obstructive sleep apnea: A systematic review and meta-analysis. *Sleep Medicne Reviews*, *45*, 1–17. https://doi.org/10.1016/j.smrv.2019.01.004

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Zheng, B., Yu, C., Lv, J., Guo, Y., Bian, Z., Zhou, M., Yang, L., Chen, Y., Li, X., Ning, F., Chen, J., Chen, Z., & Li, L. (2019). Insomnia symptoms and risk of cardiovascular diseqases among 0.5 million adults: A 10-year cohort. *Neurology*, *93*, e2110–e2120.

Zheng, Y., Lv, T., Wu, J., & Lyu, Y. (2022). Trazodone changed the polysomnographic sleep architecture in insomnia disorder: A systematic review and meta-analysis. *Scientific Reports*, *12*, 14453.

Zhou, F. C., Yang, Y., Wang, Y. Y., Rao, W. W., Zhang, S. F., Zeng, L. N., Zheng, W., Ng, C. H., Ungvari, G. S., Zhang, L., & Xiang, Y. T. (2020). Cognitive behavioural therapy for insomnia monotherapy in patients with medical or psychiatric comorbidities: A meta-analysis of randomized controlled trials. *Psychiatric Quarterly*, *91*, 1209–1224.

Zhou, H., Zhao, Y., Peng, W., Han, W., Wang, D., Wang, Z., Ren, X., Pang, G., Lin, Q., & Wang, X. (2022). Efficacy and safety of Wuling capsule for insomnia disorder: A systematic review and meta-analysis of randomized controlled trials. *Sleep Medicine*, *93*, 1–14.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Riemann, D., Espie, C. A., Altena, E., Arnardottir, E. S., Baglioni, C., Bassetti, C. L. A., Bastien, C., Berzina, N., Bjorvatn, B., Dikeos, D., Dolenc Groselj, L., Ellis, J. G., Garcia-Borreguero, D., Geoffroy, P. A., Gjerstad, M., Gonçalves, M., Hertenstein, E., Hoedlmoser, K., Hion, T., … Spiegelhalder, K. (2023). The European Insomnia Guideline: An update on the diagnosis and treatment of insomnia 2023. *Journal of Sleep Research*, *32*(6), e14035. https://doi.org/10.1111/jsr.14035

13652869, 2023, 6, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/jsr.14035 by Cochrane Portugal, Wiley Online Library on [07/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License